

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
8 April 2004 (08.04.2004)

PCT

(10) International Publication Number
WO 2004/028444 A2

- (51) International Patent Classification⁷: A61K
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (21) International Application Number: PCT/US2002/025348
- (22) International Filing Date: 9 August 2002 (09.08.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (71) Applicant: SCANTIBODIES LABORATORY, INC. [US/US]; 9336 Abraham Way, Santee, CA 92071 (US).
- (72) Inventor: CANTOR, Thomas, L.; 11149 Shining Light Way, El Cajon, CA 92020 (US).
- (74) Agents: CHEN, Peng et al.; Morrison & Foerster LLP, 3811 Valley Centre Drive, Suite 500, San Diego, CA 92130-2332 (US).
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PARATHYROID HORMONE ANTAGONISTS AND USES THEREOF



(57) Abstract: The present invention relates to parathyroid hormone (PTH) antagonists. More particularly, the present invention provides for pharmaceutical compositions, kits and combinations comprising the PTH antagonist. The present invention also provides for methods for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, e.g., calcium, loss or for preventing, treating or delaying the effect of a PTH agonist using the PTH antagonist. The present invention further provides for methods for identifying a subject having or at risk of having osteoporosis or decreased bone density, or for identifying a subject in need of PTH antagonist treatment, or for monitoring a subject undergoing treatment for osteoporosis or decreased bone density, by determining and/or monitoring PTH antagonist level or a comparative value between PTH agonist and PTH antagonist. The present invention further provides for methods for identifying an agent suitable for preventing, treating or delaying osteoporosis by identifying a compound that enhances the PTH antagonist activity.

Best Available Copy

WO 2004/028444 A2

PARATHYROID HORMONE ANTAGONISTS AND USES THEREOF

[0001] The present application is a continuation-in-part application of the following U.S. patent applications: 1) Serial No. 09/323,606, filed June 2, 1999;
5 2) Serial No. 09/928,047, filed August 10, 2001, which claims priority benefit of U.S. provisional application Serial No. 60/224,446, filed August 10, 2000; and 3) Serial No. 09/636,530, filed August 10, 2001, which claims priority benefit of U.S. provisional application Serial No. 60/224,447, filed August 10, 2000. The disclosure of the above-referenced applications is incorporated by reference
10 herein in their entireties.

Technical Field

[0002] The present invention relates to parathyroid hormone (PTH) antagonists. More particularly, the present invention provides for pharmaceutical compositions, kits and combinations comprising the PTH antagonist. The present
15 invention also provides for methods for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, e.g., calcium, loss or for preventing, treating or delaying the effect of a PTH agonist using the PTH antagonist. The present invention further provides for methods for identifying a subject having or at risk of having osteoporosis or decreased bone density, or for
20 identifying a subject in need of PTH antagonist treatment, or for monitoring a subject undergoing treatment for osteoporosis or decreased bone density, by determining and/or monitoring PTH antagonist level or a comparative value between PTH agonist and PTH antagonist. The present invention further provides for methods for identifying an agent suitable for preventing, treating or delaying
25 osteoporosis by identifying a compound that enhances the PTH antagonist activity.

BACKGROUND OF THE INVENTION

[0003] Calcium plays an indispensable role in cell permeability, the formation of bones and teeth, blood coagulation, transmission of nerve impulse,

and normal muscle contraction. The concentration of calcium ions in the blood is, along with calcitriol and calcitonin, regulated mainly by parathyroid hormone (PTH). Extracellular calcium levels are directly affected by PTH through calcium uptake in kidney tubule cells and calcium transport to or from bone. Although calcium intake and excretion may vary, PTH serves through a feedback mechanism to maintain a steady concentration of calcium in cells and surrounding fluids. When serum calcium lowers, the parathyroid glands secrete PTH, affecting the release of stored calcium. When serum calcium increases, stored calcium release is retarded through lowered secretions of PTH.

[0004] The complete or whole form of human PTH, (hPTH), is a unique 84 amino acid peptide (SEQ ID NO:1), as is shown in FIGURE 1. Researchers have found that this peptide has an anabolic effect on bone that involves a domain for protein kinase C activation (amino acid residues 28 to 34) as well as a domain for adenylate cyclase activation (amino acid residues 1 to 7). However, various catabolic forms of clipped or fragmented PTH peptides also are found in circulation, most likely formed by intraglandular or peripheral metabolism. For example, hPTH can be cleaved between amino acids 34 and 35 to produce a (1-34) PTH N-terminal fragment and a (35-84) PTH C-terminal fragment. Likewise, clipping can occur between either amino acids 36 and 37 or 37 and 38. Recently, a large PTH fragment referred to as "non-(1-84) PTH" has been disclosed which is clipped closer to the N-terminal end of PTH. (See R. LePage et al., *Clin. Chem.*, 44:805-810 (1998)).

[0005] The cleaved fragments of PTH vary in both biological activity and metabolic clearance rate from the circulation. For example, the N-terminal human PTH₁₋₃₄ (hPTH₁₋₃₄) fragment has PTH agonist properties, but is rapidly removed from circulation. A daily subcutaneous administration of hPTH to patients with idiopathic osteoporosis has been shown to substantially increase their iliac trabecular bone volume. (See Podbesek et al., *Endocrinology*, 112:1000-1006 (1983)).

[0006] PTH plays a role in the course of disease in a patient with chronic renal failure. Renal osteodystrophy (RO) is a complex skeletal disease

comprising osteitis fibrosa cystica (caused by excess PTH), osteomalacia resulting in mineralized bone matrix (caused by vitamin D deficiency), extraskeletal calcification/ossification (caused by abnormal calcium and phosphorus metabolism), and adynamic bone disease (contributed to by PTH suppression).

- 5 Chronic renal failure patients can develop RO. Failing kidneys increase serum phosphorus (hyperphosphoremia) and decrease 1,25-dihydroxyvitamin D (1,25-D) production by the kidney. The former results in secondary hyperparathyroidism from decreased gastrointestinal calcium absorption and osteitis fibrosa cystica from increased PTH in indirect response to an increase in
10 serum phosphorus. The later causes hypocalcemia and osteomalacia. With the onset of secondary hyperparathyroidism, the parathyroid gland becomes less responsive to its hormonal regulators because of decreased expression of its calcium and vitamin D receptors. Serum calcium drops. RO can lead to digital gangrene, bone pain, bone fractures, and muscle weakness.

- 15 [0007] To treat secondary hyperparathyroidism, patients are given calcium and vitamin D replacement. Vitamin D analogues, such as calcitriol, stimulate intestinal calcium transport, calcium absorption in bone and calcium tubular reabsorption in kidneys. Such therapy has its dangers. Serum calcium levels must be carefully monitored. Too much dosage can induce hypercalcemia or
20 hypercalciuria. Moreover, very serious consequences occur from calcium and phosphorus mismanagement from direct and indirect PTH suppression therapy. Soft tissue calcification results in a five to fifteen times higher incidence of myocardial infarction among end stage renal dialysis patients as compared to age matched diabetes patients. The secondarily hyperplastic parathyroid glands
25 escape PTH control over calcium, a condition referred to as tertiary hyperparathyroidism.

- [0008] Another treatment proposed for patients with excess PTH is to administer parathyroid hormone analogues which inhibit the biological activity of PTH. U.S. Patent Nos. 5,093,233 and 4,968,669 disclose N-terminal PTH
30 analogues (PTH₇₋₃₄ and PTH₈₋₃₄), having substitutions at the 8, 12, 18, and/or 34 amino acid positions. These analogs bind to PTH cell surface receptors but do not

stimulate a change in the second messenger concentration, *i.e.*, act as a hormone for calcium ion concentration. PTH activity can also be inhibited by unsubstituted PTH fragments, namely PTH₁₋₃₄ or PTH₇₋₃₄, however, these fragments are so weak in their antagonist properties that they do not have practical or beneficial significance.

[0009] Osteoporosis is the most common form of metabolic bone disease and may be considered the symptomatic, fracture stage of bone loss (osteopenia). Although osteoporosis may occur secondary to a number of underlying diseases, 90% of all cases appear to be idiopathic. Postmenopausal women are particularly at risk for idiopathic osteoporosis (postmenopausal or Type I osteoporosis). Another high risk group for idiopathic osteoporosis is the elderly of either sex (senile or Type II osteoporosis). Osteoporosis has also been related to corticosteroid use, immobilization or extended bed rest, alcoholism, diabetes, gonadotoxic chemotherapy, hyperprolactinemia, anorexia nervosa, primary and secondary amenorrhea, and oophorectomy.

[0010] In the various forms of osteoporosis, mechanical failure bone fractures frequently occur which are the result of bone loss. Postmenopausal osteoporosis is characterized by fractures of the wrist and spine, while femoral neck fractures seem to be the dominant feature of senile osteoporosis.

[0011] Bone loss in osteoporotics is believed to involve an imbalance in the process by which the skeleton renews itself. This process has been termed bone remodeling. It occurs in a series of discrete pockets of activity. These pockets appear spontaneously within the bone matrix on a given bone surface as a site of bone resorption. Osteoclasts (bone dissolving or resorbing cells) are responsible for the resorption of a portion of bone of generally constant dimension. Resorption is followed by the appearance of osteoblasts (bone forming cells) that refill the cavity left by the osteoclasts with new bone.

[0012] In a healthy adult subject, the rate at which osteoclasts and osteoblasts are formed is such that bone formation and bone resorption are in balance constituting an optimal bone turnover rate. However, in osteoporotics an imbalance in the bone remodeling process develops which results in bone being

lost at a rate faster than it is being made. Although this imbalance occurs to some extent in most individuals as they age, it is much more severe and occurs at a younger age in postmenopausal osteoporotics or following oophorectomy.

[0013] There have been many attempts to treat osteoporosis with the goal of either slowing further bone loss or, more desirably, producing a net gain in bone mass. Certain agents, such as estrogen and the bisphosphonates, appear to slow further bone loss in osteoporotics. Agents which slow bone loss, because of the different durations of bone resorption and formation, may appear to increase bone mass (on the order of 3% to 7%). However, this apparent increase is limited in time, not progressive, and is due to a decrease in "remodeling space." In addition, because of the close coupling between resorption and formation, impeding bone resorption also ultimately impedes bone formation.

[0014] Another class of agents investigated to combat the onset of osteoporosis encompasses PTH and PTH analogues. (See e.g., U.S. Patent No. 6,051,686). The theory behind the use of such compounds is to use the body's natural protein receptor binding process to counter a greater removal of calcium from bone than resorption of calcium. Unfortunately, such proposed treatments have had adverse effects, including hypercalcemia (elevated serum calcium) and the formation of osteosarcomas.

[0015] There exists a need in the art for more compositions and methods for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, e.g., calcium, loss or for preventing, treating or delaying the effect of a PTH agonist. The present invention addresses this and other related needs in the art.

BRIEF SUMMARY OF THE INVENTION

[0016] In one aspect, the present invention is directed to a parathyroid hormone (PTH) antagonist, which PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₃₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue

of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues.

5 Preferably, the PTH antagonist is in the form of a pharmaceutical composition, which pharmaceutical composition comprises an effective amount of the PTH antagonist and a pharmaceutically acceptable carrier or excipient. Kits and combinations comprising the PTH antagonist are also provided herein. In a specific embodiment, the PTH antagonist is not PTH₇₋₈₄.

10 [0017] In another aspect, the present invention is directed to a method for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH)
15 antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue
20 of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby said disease or disorder associated with excessive bone mineral, *e.g.*,
25 calcium, loss is prevented, treated or delayed. In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

[0018] In still another aspect, the present invention is directed to a method for preventing, treating or delaying the effect of a PTH agonist in a mammal, which method comprises administering to a mammal, to which such prevention,
30 treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or

antagonizing function of said PTH antagonist wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby the effect of said PTH agonist is prevented, treated or delayed.

In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

[0019] In yet another aspect, the present invention is directed to a method for identifying a subject having or at risk of having osteoporosis or decreased bone density, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist as having or at risk of having osteoporosis or decreased bone density. In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

[0020] In yet another aspect, the present invention is directed to a method for identifying a subject in need of parathyroid hormone (PTH) antagonist treatment, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist as in need of parathyroid hormone (PTH) antagonist treatment. In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

[0021] In yet another aspect, the present invention is directed to a method for monitoring a subject undergoing treatment for osteoporosis or decreased bone density, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a

subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist. In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

[0022] In yet another aspect, the present invention is directed to a method for identifying an agent suitable for preventing, treating or delaying osteoporosis, which method comprises: a) measuring PTH antagonist activity in the presence and absence of a test substance (or candidate compound); and b) identifying a test substance (or candidate compound) that enhances said PTH antagonist activity as an agent suitable for preventing, treating or delaying osteoporosis. In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

[0023] In yet another aspect, the present invention is directed to a pharmaceutical composition, which pharmaceutical composition comprises an effective amount of a parathyroid hormone (PTH) antagonist and a pharmaceutically acceptable carrier or excipient, wherein said PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set forth in SEQ ID NO:2 (svseiqlmh nlghlssle rvewlrkkqlq dvhnfvalga sivhrdgssq rprkkednvl veshqkslge adkaavdvli kakpq), dog PTH having an amino acid sequence set forth in SEQ ID NO:3 (svseiqlmh nlghlssme rvewlrkkqlq dvhnfvalga piahrdgssq rplkkednvl veshqkslge adkadvdvlt kaksq), bovine PTH having an amino acid sequence set forth in SEQ ID NO:4 (avseiqlmh nlghlssme rvewlrkkqlq dvhnfvalga siayrdgssq rprkkednvl veshqkslge adkadvdvli kakpq), rat PTH having an amino acid sequence set forth in SEQ ID NO:5 (avseiqlmh nlghlasve rmqwrkkqlq dvhnfvslgv qmaaregsyq rptkkeenvl vdgnskslge gdkadvdvltv kaksq), or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6 (svsemqlmh nlgehrhtve rqdwlqmklq dvhsaledar tqrpmkedi vlgeimrrl lpehlaavq kksidldkay mnvlfktkp), or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a

minimal length of three amino acid residues. In a specific embodiment, the PTH antagonist used in the present pharmaceutical composition is not PTH₇₋₈₄.

[0024] In yet another aspect, the present invention is directed to a method for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set forth in SEQ ID NO:2, dog PTH having an amino acid sequence set forth in SEQ ID NO:3, bovine PTH having an amino acid sequence set forth in SEQ ID NO:4, rat PTH having an amino acid sequence set forth in SEQ ID NO:5 or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6, or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby said disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss is prevented, treated or delayed. In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

[0025] In yet another aspect, the present invention is directed to a method for preventing, treating or delaying the effect of a PTH agonist in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set forth in SEQ ID NO:2, dog PTH having an amino acid sequence set forth in SEQ

ID NO:3, bovine PTH having an amino acid sequence set forth in SEQ ID NO:4, rat PTH having an amino acid sequence set forth in SEQ ID NO:5, or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6, or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby the effect of said PTH agonist is prevented, treated or delayed. In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

[0026] In yet another aspect, the present invention is directed to a method for preventing, treating or delaying bone metastasis in a human, which method comprises administering to a human, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby said bone metastasis is prevented, treated or delayed. In a specific embodiment, the PTH antagonist used in the present method is not PTH₇₋₈₄.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWING(S)

[0027] Figure 1 is a diagrammatic view of human PTH (hPTH).

[0028] Figure 2 illustrates comparison of the recognition of hPTH 1-84 and hPTH 7-84 by the Nichols I-PTH assay. The Nichols I-PTH assay does not differentiate between hPTH 1-84 (solid line) and hPTH 7-84 (dashed line).

[0029] Figure 3 illustrates comparison of the recognition of hPTH 1-84 and hPTH 7-84 by the Whole PTH assay. Unlike the Nichols I-PTH assay, the Whole PTH assay does discriminate between hPTH 1-84 (solid line) and hPTH 7-84 (dashed line). Concentrations of hPTH 7-84 as high as 10,000 pg were undetectable.

[0030] Figure 4 illustrates comparison of the effect of hPTH 1-84 or hPTH 7-84 on cAMP production in ROS 17.2 cells. Unlike hPTH 7-84, hPTH 1-84 increased cAMP production in a dose-dependent manner. cAMP increased from 18.1 ± 1.2 to 738 ± 4.1 nmol/well after treatment with 10^{-8} mol/L hPTH 1-84. The same concentration of hPTH 7-84 had no effect.

[0031] Figure 5 illustrates comparison of the calcemic effects of PTH isoforms. Parathyroidectomized (PTX) rats fed a 0.02% calcium diet show a significant increase in plasma calcium after treatment with hPTH 1-84. In contrast, hPTH 7-84 produced a slight but significant decrease in plasma calcium. When both peptides were given together in a 1:1 molar ratio, the calcemic response induced by hPTH 1-84 was reduced by 94% ($P < 0.001$). Symbols are: (●) 1-84, N = 9; (■) 1-84 + 7-84, N = 6; (▲) Control, N = 5; (◆) 7-84, N = 5.

[0032] Figure 6 illustrates comparison of the effects of hPTH 1-84 or hPTH 1-84 plus hPTH 7-84 on (A) glomerular filtration rate (GFR) and (B) fractional excretion of phosphorus (FE_{po4}). Control and treatment periods are denoted by open and closed bars, respectively. The phosphaturia induced by hPTH 1-84 was decreased by 50.2% ($P < 0.05$) when animals were treated simultaneously with 7-84 PTH, despite a significant increase in GFR ($P < 0.005$).

[0033] Figure 7 illustrates comparison of PTH values in plasma from uremic patients using the Nichols "intact" PTH assay (■) versus the Whole PTH assay (●). Plasma PTH values are uniformly higher when measured with the Nichols "intact" PTH assay than with the Whole PTH assay. The median PTH values were 523 vs. 344 pg/mL, respectively ($P < 0.001$).

[0034] Figure 8 illustrates regression analysis of plasma PTH measured by Nichols I-PTH and Whole PTH assay in uremic patients ($r=0.97$; $P<0.001$).

[0035] Figure 9 illustrates effects of plasma calcium on PTH degradation in dialysis patients. The percentage of non-(1-84) PTH fragment (likely hPTH 7-84) correlates positively with plasma calcium ($P<0.02$) ($r=0.638$; $P=0.0025$; $N=20$).

[0036] Figure 10 illustrates comparison of plasma PTH levels in renal transplant patients using Nichols I-PTH and Whole PTH assays. PTH values are higher when measured with the Nichols I-PTH assay ($P>0.005$).

[0037] Figure 11 illustrates intracellular PTH content on parathyroid glands from uremic patients. The $41.8 \pm 3.2\%$ of the total PTH, measured by the I-Nichols assay (expressed as 100%), represents the non-(1-84) PTH fragment "likely" hPTH 7-84 (\square). the 1-84 PTH molecule was measured with the Whole PTH assay (\blacksquare).

DETAILED DESCRIPTION OF THE INVENTION

[0038] For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the subsections that follow.

A. Definitions

[0039] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this invention belongs. All patents, applications, published applications and other publications referred to herein are incorporated by reference in their entirety. If a definition set forth in this section is contrary to or otherwise inconsistent with a definition set forth in the patents, applications, published applications and other publications that are herein incorporated by reference, the definition set forth in this section prevails over the definition that is incorporated herein by reference.

[0040] As used herein, "a" or "an" means "at least one" or "one or more."

[0041] As used herein, "parathyroid hormone (PTH) agonist" refers to the complete molecule of PTH or a fragment or derivative thereof that stimulates osteoclasts formation and bone turnover to increase blood calcium levels. Other names of PTH include parathormone and parathyrin. For purposes herein, the name "parathyroid hormone (PTH)" is used herein, although all other names are contemplated. It is intended to encompass PTH agonist with conservative amino acid substitutions that do not substantially alter its activity. Suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson *et al.*, *Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacin/Cummings Pub. co., p.224).

[0042] As used herein, "parathyroid hormone (PTH) antagonist" refers to a PTH fragment or derivative that counters the effect of a PTH agonist. It is intended to encompass PTH antagonist with conservative amino acid substitutions that do not substantially alter its activity. Suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson *et al.*, *Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacin/Cummings Pub. co., p.224).

[0043] As used herein, a "functional derivative or fragment" of PTH agonist or PTH antagonist refers to a derivative or fragment of PTH that still substantially retains its function as a PTH agonist or PTH antagonist. Normally, the derivative or fragment retains at least 50% of its PTH agonist or PTH antagonist activity. Preferably, the derivative or fragment retains at least 60%, 70%, 80%, 90%, 95%, 99% and 100% of its PTH agonist or PTH antagonist activity. It is also possible that a functional derivative or fragment of PTH agonist

or PTH antagonist has higher PTH agonist or PTH antagonist activity than a parent molecule from which the functional derivative or fragment is derived from.

[0044] As used herein, "an agent that enhances production of said PTH antagonist" refers to a substance that increases transcription and/or translation of a PTH antagonist, or a substance that increases post-translational modification and/or cellular trafficking of a PTH antagonist precursor, or a substance, *e.g.*, a protease, that cleaves or a substance that enhances the cleavage a PTH protein to generate a PTH antagonist, or a substance that prolongs half-life of a PTH antagonist.

[0045] As used herein, an "agent that enhances antagonizing function of said PTH antagonist" refers to a substance that increases potency of the PTH antagonist, or a substance that increases sensitivity of PTH antagonist's natural ligand in a PTH antagonist signaling pathway, or a substance that decreases potency of a PTH agonist.

[0046] As used herein, "an abnormal PTH agonist level" refers to: 1) a PTH agonist level in a individual mammal, *e.g.*, human, that is statistically higher or lower than a PTH agonist level in the same individual mammal in a healthy state; 2) a PTH agonist level in a individual mammal, *e.g.*, human, that is statistically higher or lower than a PTH agonist level in another comparable individual mammal in a healthy state; or 3) a PTH agonist level in a individual mammal, *e.g.*, human, that is statistically higher or lower than a mean or average PTH agonist level of comparable healthy population. The difference between the abnormal PTH agonist level and the normal PTH agonist level must be statistically significant so that the difference or the abnormal PTH agonist level can be used in prognosis, diagnosis or treatment monitoring.

[0047] As used herein, "an abnormal PTH antagonist level" refers to: 1) a PTH antagonist level in a individual mammal, *e.g.*, human, that is statistically higher or lower than a PTH antagonist level in the same individual mammal in a healthy state; 2) a PTH antagonist level in a individual mammal, *e.g.*, human, that is statistically higher or lower than a PTH antagonist level in another comparable individual mammal in a healthy state; or 3) a PTH antagonist level in a individual

mammal, *e.g.*, human, that is statistically higher or lower than a mean or average PTH antagonist level of comparable healthy population. The difference between the abnormal PTH antagonist level and the normal PTH antagonist level must be statistically significant so that the difference or the abnormal PTH antagonist level can be used in prognosis, diagnosis or treatment monitoring.

[0048] As used herein, "an abnormal comparative value between PTH agonist and PTH antagonist" refers to: 1) a comparative value between PTH agonist and PTH antagonist in a individual mammal, *e.g.*, human, that is statistically higher or lower than such a comparative value in the same individual mammal in a healthy state; 2) a comparative value between PTH agonist and PTH antagonist in a individual mammal, *e.g.*, human, that is statistically higher or lower than such a comparative value in another comparable individual mammal in a healthy state; or 3) a comparative value between PTH agonist and PTH antagonist in a individual mammal, *e.g.*, human, that is statistically higher or lower than a mean or average comparative value of comparable healthy population. The difference between the abnormal comparative value and the normal comparative value must be statistically significant so that the difference or the abnormal comparative value between PTH agonist and PTH antagonist can be used in prognosis, diagnosis or treatment monitoring. The comparative value between PTH agonist and PTH antagonist can take any suitable form. For example, the comparative value can be a ratio, *e.g.*, PTH agonist/PTH antagonist, PTH antagonist/PTH agonist, PTH agonist/the sum of PTH agonist and PTH antagonist, or PTH antagonist/the sum of PTH agonist and PTH antagonist, etc. In another example, the comparative value can be a subtraction value, *e.g.*, PTH agonist-PTH antagonist, PTH antagonist-PTH agonist, etc. The above examples are for illustration only and are not intended to be an exhaustive list of all possible formats for measuring the comparative value between PTH agonist and PTH antagonist. Other suitable formats are readily apparent to skilled artisans and can be used.

[0049] As used herein, "a disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss" refers to any disease or disorder associated with or caused by net bone loss.

5 [0050] As used herein, "an agent suitable for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss" does not encompass a PTH antagonist.

[0051] As used herein, "an effective amount of a compound for treating a particular disease" is an amount that is sufficient to ameliorate, or in some manner reduce the symptoms associated with the disease. Such amount may be
10 administered as a single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but, typically, is administered in order to ameliorate the symptoms of the disease. Repeated administration may be required to achieve the desired amelioration of symptoms.

[0052] As used herein, "treatment" means any manner in which the
15 symptoms of a condition, disorder or disease are ameliorated or otherwise beneficially altered. Treatment also encompasses any pharmaceutical use of the compositions herein.

[0053] As used herein, "amelioration" of the symptoms of a particular disorder by administration of a particular pharmaceutical composition refers to
20 any lessening, whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the composition.

[0054] As used herein, "production by recombinant means" refers to production methods that use recombinant nucleic acid methods that rely on well known methods of molecular biology for expressing proteins encoded by cloned
25 nucleic acids.

[0055] As used herein, "complementary" when referring to two nucleic acid molecules, means that the two sequences of nucleotides are capable of hybridizing, preferably with less than 25%, more preferably with less than 15%, even more preferably with less than 5%, most preferably with no mismatches
30 between opposed nucleotides. Preferably the two molecules will hybridize under conditions of high stringency.

[0056] As used herein: "stringency of hybridization" in determining percentage mismatch is as follows:

[0057] 1) high stringency: 0.1 x SSPE, 0.1% SDS, 65°C;

[0058] 2) medium stringency: 0.2 x SSPE, 0.1% SDS, 50°C (also referred to as moderate stringency); and

[0059] 3) low stringency: 1.0 x SSPE, 0.1% SDS, 50°C.

[0060] It is understood that equivalent stringencies may be achieved using alternative buffers, salts and temperatures.

[0061] As used herein, "vector (or plasmid)" refers to discrete elements that are used to introduce heterologous DNA into cells for either expression or replication thereof. Selection and use of such vehicles are well known within the skill of the artisan. An expression vector includes vectors capable of expressing DNA's that are operatively linked with regulatory sequences, such as promoter regions, that are capable of effecting expression of such DNA fragments. Thus, an expression vector refers to a recombinant DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other vector that, upon introduction into an appropriate host cell, results in expression of the cloned DNA. Appropriate expression vectors are well known to those of skill in the art and include those that are replicable in eucaryotic cells and/or prokaryotic cells and those that remain episomal or those which integrate into the host cell genome.

[0062] As used herein, "a promoter region or promoter element" refers to a segment of DNA or RNA that controls transcription of the DNA or RNA to which it is operatively linked. The promoter region includes specific sequences that are sufficient for RNA polymerase recognition, binding and transcription initiation. This portion of the promoter region is referred to as the promoter. In addition, the promoter region includes sequences that modulate this recognition, binding and transcription initiation activity of RNA polymerase. These sequences may be *cis* acting or may be responsive to *trans* acting factors. Promoters, depending upon the nature of the regulation, may be constitutive or regulated.

Exemplary promoters contemplated for use in prokaryotes include the bacteriophage T7 and T3 promoters, and the like.

[0063] As used herein, "operatively linked or operationally associated" refers to the functional relationship of DNA with regulatory and effector sequences of nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other signal sequences. For example, operative linkage of DNA to a promoter refers to the physical and functional relationship between the DNA and the promoter such that the transcription of such DNA is initiated from the promoter by an RNA polymerase that specifically recognizes, binds to and transcribes the DNA. In order to optimize expression and/or *in vitro* transcription, it may be necessary to remove, add or alter 5' untranslated portions of the clones to eliminate extra, potential inappropriate alternative translation initiation (*i.e.*, start) codons or other sequences that may interfere with or reduce expression, either at the level of transcription or translation. Alternatively, consensus ribosome binding sites (see, *e.g.*, Kozak, *J. Biol. Chem.*, 266:19867-19870 (1991)) can be inserted immediately 5' of the start codon and may enhance expression. The desirability of (or need for) such modification may be empirically determined.

[0064] As used herein, "pharmaceutically acceptable salts, esters or other derivatives" include any salts, esters or derivatives that may be readily prepared by those of skill in this art using known methods for such derivatization and that produce compounds that may be administered to animals or humans without substantial toxic effects and that either are pharmaceutically active or are prodrugs.

[0065] As used herein, a "prodrug" is a compound that, upon *in vivo* administration, is metabolized or otherwise converted to the biologically, pharmaceutically or therapeutically active form of the compound. To produce a prodrug, the pharmaceutically active compound is modified such that the active compound will be regenerated by metabolic processes. The prodrug may be designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other

characteristics or properties of a drug. By virtue of knowledge of pharmacodynamic processes and drug metabolism *in vivo*, those of skill in this art, once a pharmaceutically active compound is known, can design prodrugs of the compound (see, e.g., Nogrady (1985) Medicinal Chemistry A Biochemical Approach, Oxford University Press, New York, pages 388-392).

5 [0066] As used herein, "test substance (or candidate compound)" refers to a chemically defined compound (e.g., organic molecules, inorganic molecules, organic/inorganic molecules, proteins, peptides, nucleic acids, oligonucleotides, lipids, polysaccharides, saccharides, or hybrids among these molecules such as
10 glycoproteins, etc.) or mixtures of compounds (e.g., a library of test compounds, natural extracts or culture supernatants, etc.) whose effect on PTH antagonist is determined by the disclosed and/or claimed methods herein.

[0067] As used herein, high-throughput screening (HTS) refers to processes that test a large number of samples, such as samples of diverse chemical
15 structures against disease targets to identify "hits" (see, e.g., Broach, *et al.*, High throughput screening for drug discovery, *Nature*, 384:14-16 (1996); Janzen, *et al.*, High throughput screening as a discovery tool in the pharmaceutical industry, *Lab Robotics Automation*: 8261-265 (1996); Fernandes, P.B., Letter from the society president, *J. Biomol. Screening*, 2:1 (1997); Burbaum, *et al.*, New technologies for
20 high-throughput screening, *Curr. Opin. Chem. Biol.*, 1:72-78 (1997)). HTS operations are highly automated and computerized to handle sample preparation, assay procedures and the subsequent processing of large volumes of data.

[0068] As used herein, "disease or disorder" refers to a pathological condition in an organism resulting from, e.g., infection or genetic defect, and
25 characterized by identifiable symptoms.

B. PTH antagonists and pharmaceutical composition, kits and combinations comprising the same

30 [0069] In one aspect, the present invention is directed to a parathyroid hormone (PTH) antagonist, which PTH antagonist comprises a contiguous portion

of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues.

Preferably, the PTH antagonist is in the form of a pharmaceutical composition, which pharmaceutical composition comprises an effective amount of the PTH antagonist and a pharmaceutically acceptable carrier or excipient.

[0070] The N-terminal amino acid residue of the PTH antagonist can start at any position spanning position 2 through position 33 of said PTH₁₋₈₄. For example, the N-terminal amino acid residue of the PTH antagonist can start at position 2 of the PTH₁₋₈₄. The C-terminal amino acid residue of said PTH antagonist can end at any position spanning position 35 through position 84 of said PTH₁₋₈₄. For example, the C-terminal amino acid residue of the PTH antagonist can end at position 84 of the PTH₁₋₈₄.

[0071] In a specific embodiment, the PTH antagonist is a protein or a peptide, or a nucleic acid encoding said protein or peptide, selected from the group consisting of PTH₂₋₈₄, PTH₃₋₈₄, PTH₄₋₈₄, PTH₅₋₈₄, PTH₆₋₈₄, PTH₇₋₈₄, PTH₈₋₈₄, PTH₉₋₈₄, PTH₁₀₋₈₄, PTH₁₁₋₈₄, PTH₁₂₋₈₄, PTH₁₃₋₈₄, PTH₁₄₋₈₄, PTH₁₅₋₈₄, PTH₁₆₋₈₄, PTH₁₇₋₈₄, PTH₁₈₋₈₄, PTH₁₉₋₈₄, PTH₂₀₋₈₄, PTH₂₁₋₈₄, PTH₂₂₋₈₄, PTH₂₃₋₈₄, PTH₂₄₋₈₄, PTH₂₅₋₈₄, PTH₂₆₋₈₄, PTH₂₇₋₈₄, PTH₂₈₋₈₄, PTH₂₉₋₈₄, PTH₃₀₋₈₄, PTH₃₁₋₈₄, PTH₃₂₋₈₄, and PTH₃₃₋₈₄. In another specific embodiment, the PTH antagonist is a protein or a peptide, or a nucleic acid encoding said protein or peptide, selected from the group consisting of PTH₇₋₆₉, PTH₇₋₇₀, PTH₇₋₇₁, PTH₇₋₇₂, PTH₇₋₇₃, PTH₇₋₇₄, PTH₇₋₇₅, PTH₇₋₇₆, PTH₇₋₇₇, PTH₇₋₇₈, PTH₇₋₇₉, PTH₇₋₈₀, PTH₇₋₈₁, PTH₇₋₈₂, PTH₇₋₈₃ and PTH₇₋₈₄.

[0072] The PTH antagonist can have any suitable length provided that it maintains its antagonizing activity. For example, the PTH antagonist can have a length of 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23,

24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82 or 83 amino acid residues.

[0073] The PTH antagonist can further comprise an amino acid residue

5 substitution or modification that enhances or does not decrease its antagonist activity, or an amino acid residue substitution or modification that stabilizes the PTH antagonist. For example, the PTH antagonist can further comprise the following amino acid residue substitution or modification: His₂₅, His₂₆, Leu₂₇, (U.S. Patent No. 5,382,658); Tyr₃₄, D-Trp₁₂, Nle_{8,18}, desamino(Nle_{8,18}), Lys₁₃
10 modified in the epsilon-amino acid group by N,N-diisobutyl or 3-phenylpropanoyl (U.S. Patent No. 5,093,233); Gly₁₂ substituted by D-Trp, L-Trp, L- or D- α - or β -naphthylalanine, or D- or L- α -MeTrp (U.S. Patent No. 4,968,669); the amino acid residue at positions 7, 11, 23, 24, 27, 28, or 31 being cyclohexylalanine, the amino acid residue at position 3, 16, 17, 18, 19, or 34 being α -aminoisobutyric acid, the
15 amino acid residue at position 1 being α , β -diaminopropionic acid, the amino acid residue at position 27 being homoarginine, the amino acid residue at position 31 being norleucine (U.S. Patent No. 5,723,577); each of Arg₂₅, Lys₂₆, Lys₂₇ being substituted with Ala, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Met, Phe, Pro, Ser, Thr, Trp, Tyr or Val (U.S. Patent No. 5,317,010); and a combination thereof.

20 [0074] The PTH antagonist or a pharmaceutical composition comprising the same can be formulated in any suitable formats. For example, the PTH antagonist or a pharmaceutical composition comprising the same can be formulated in a solid or a liquid dosage form. In another example, the PTH antagonist or a pharmaceutical composition comprising the same can be
25 formulated for oral, parenteral, intranasal, topical, or injectable administration, e.g., intracavernous injection, subcutaneous injection, intravenous injection, intramuscular injection and intradermal injection.

[0075] If a nucleic acid encoding a PTH antagonist is used, the nucleic acid can be in any suitable forms, e.g., DNA, RNA, PNA or a combination
30 thereof. In a specific example, the nucleic acid can be comprised in a gene

therapy vector, e.g., an adenovirus associated vector, a retroviral vector, an adenovirus vector, and a lentivirus vector.

- [0076] PTH antagonist proteins or functional derivatives or fragments thereof, or a nucleic acid encoding PTH antagonist, or functional derivatives or fragments thereof, can be prepared by any methods known in the art, e.g., synthetic methods, recombinant methods or a combination thereof (*See generally*, Ausubel (Ed.) *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (2000)).
- [0077] PTH antagonist proteins or functional derivatives or fragments thereof, or a nucleic acid encoding PTH antagonist, or functional derivatives or fragments thereof, derived from known PTH encoding gene can also be used, e.g., AF251060 (Homo sapiens parathyroid hormone); AF130257 (Macaca fascicularis parathyroid hormone); AH007117 (Mus musculus parathyroid hormone); AF066075 (Mus musculus parathyroid hormone); U15662 (Canis familiaris parathyroid hormone, Rosol et al., *Gene*, 160(2):241-243 (1995)); M26143 (Synthetic human parathyroid hormone, Sung et al., *Biochem. Cell Biol.*, 64(2):133-138 (1986)); L19475 (rat parathyroid hormone, Pausova et al., *Genomics*, 20(1):20-26 (1994)); U17418 (human parathyroid hormone); K01938 (Bovine parathyroid hormone, Weaver et al., *Gene*, 28(3):319-329 (1984)); M25082 (Bovine parathyroid hormone, Weaver et al., *Mol. Cell. Endocrinol.*, 28(3):411-424 (1982)); P41593 (mouse parathyroid hormone, Karperien et al., *Mech. Dev.*, 47(1):29-42 (1994)); (rat parathyroid hormone, Heinrich et al., *J. Biol. Chem.*, 259(5):3320-3329 (1984)); P50133 (porcine parathyroid hormone, Smith et al., *Biochim. Biophys. Acta*, 1307(3):339-347 (1996)); P01270 (human preproparathyroid hormone, Vasicek et al., *Proc. Natl. Acad. Sci. U.S.A.*, 80(8):2127-2131 (1983)); P01268 (bovine preproparathyroid hormone, Kronenberg et al., *Proc. Natl. Acad. Sci. U.S.A.*, 76(10):4981-4985 (1979)); PTHU (parathyroid hormone precursor - human, Brewer et al., *Proc. Natl. Acad. Sci. U.S.A.*, 69(12):3585-3588 (1972)); PTBO (parathyroid hormone precursor - bovine, Brewer and Ronan, *Proc. Natl. Acad. Sci. U.S.A.*, 67(4):1862-1869

(1970); P01269 (preparathyroid hormone from pig and rat, Schmelzer et al., *Nucleic Acids Res.*, 15(16):6740 (1987)).

[0078] In another aspect, the present invention is directed to a kit, which kit comprises a PTH antagonist or a pharmaceutical composition comprising the same in a container and an instruction for using the pharmaceutical composition in preventing, treating or delaying a disease or disorder associated with excessive bone mineral, e.g., calcium, loss or for balancing the effect of excess PTH agonist. For example, the kit can be used in preventing, treating or delaying hyperparathyroidism, renal osteodystrophy, osteoporosis, parathyroid cancer, hypercalcemia, an immune disease and hypertension (See U.S. Patent No. 4,423,037).

[0079] In still another aspect, the present invention is directed to a combination, which combination comprises an effective amount of a parathyroid hormone (PTH) antagonist and an effective amount of an agent suitable for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, e.g., calcium, loss, wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues.

[0080] Any suitable PTH antagonist, including the ones described above, can be used in the present combination. Any suitable agent suitable for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, e.g., calcium, loss can be used in the present combination. For example, in treating osteoporosis, the PTH antagonist can be used in combination with known therapeutics such as calcium, bisphosphonate or vitamin D treatment.

Preferably, the present combination further comprises a pharmaceutically acceptable carrier or excipient.

[0081] In yet another aspect, the present invention is directed to a parathyroid hormone (PTH) antagonist, which PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set forth in SEQ ID NO:2, dog PTH having an amino acid sequence set forth in SEQ ID NO:3, bovine PTH having an amino acid sequence set forth in SEQ ID NO:4, rat PTH having an amino acid sequence set forth in SEQ ID NO:5, or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6, or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues. Preferably, the PTH antagonist is in the form of a pharmaceutical composition, which pharmaceutical composition comprises an effective amount of the PTH antagonist and a pharmaceutically acceptable carrier or excipient.

20 C. Methods using PTH antagonist

Methods for preventing, treating or delaying excessive bone mineral, e.g., calcium, loss

[0082] In one aspect, the present invention is directed to a method for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, e.g., calcium, loss in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋

84), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
5 c) said PTH antagonist has a minimal length of three amino acid residues, whereby said disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss is prevented, treated or delayed.

[0083] The present method can be used for preventing, treating or
10 delaying a disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss in any mammals, such as mice, rats, rabbits, cats, dogs, pigs, cows, ox, sheep, goats, horses, monkeys and other non-human primates. Preferably, the present method can be used for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss in humans.
15 For example, the human to be treated: a) is in need of increased bone density or bone healing; b) has undergone or is presently undergoing corticosteroid therapy, chemotherapy for post menopausal bone loss, radiation therapy for cancer or hormone replacement therapy; c) is immobilized or subjected to extended bed rest due to bone injury; d) suffers from alcoholism, diabetes, hyperprolactinemia,
20 anorexia nervosa, primary and secondary amenorrhea, or oophorectomy; e) is 50 years or older; f) is a female, especially a female who is 40 years or older or is in the post-menopausal stage; or g) is a male.

[0084] In a specific embodiment, the human to be treated has an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and
25 PTH antagonist. The abnormal comparative value between PTH agonist and PTH antagonist can be determined by any suitable formats, *e.g.*, as a ratio, a proportion or a subtraction value. In one example, the abnormal comparative value between PTH agonist and PTH antagonist is determined by determining and comparing at least two of the parameters selected from the group consisting of the level of the
30 PTH agonist, the PTH antagonist and the total PTH level, *i.e.*, a sum of PTH agonist and PTH antagonist. In another example, the human to be treated has a

PTH agonist/PTH antagonist ratio more than 2 and the method is used to bring the PTH agonist/PTH antagonist ratio within a range from about 1 to about 2.

[0085] The present method can be used in preventing, treating or delaying any disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss in a mammal. Exemplary diseases or disorders associated with excessive bone mineral loss, *e.g.*, calcium loss, include hyperparathyroidism, renal osteodystrophy, osteoporosis and parathyroid cancer.

[0086] Any suitable PTH antagonists can be used in the present methods. The N-terminal amino acid residue of the PTH antagonist can start at any position spanning position 2 through position 33 of said PTH₁₋₈₄. For example, the N-terminal amino acid residue of the PTH antagonist can start at position 2 of the PTH₁₋₈₄. The C-terminal amino acid residue of said PTH antagonist can end at any position spanning position 35 through position 84 of said PTH₁₋₈₄. For example, the C-terminal amino acid residue of the PTH antagonist can end at position 84 of the PTH₁₋₈₄.

[0087] In a specific embodiment, the PTH antagonist is a protein or a peptide, or a nucleic acid encoding said protein or peptide, selected from the group consisting of PTH₂₋₈₄, PTH₃₋₈₄, PTH₄₋₈₄, PTH₅₋₈₄, PTH₆₋₈₄, PTH₇₋₈₄, PTH₈₋₈₄, PTH₉₋₈₄, PTH₁₀₋₈₄, PTH₁₁₋₈₄, PTH₁₂₋₈₄, PTH₁₃₋₈₄, PTH₁₄₋₈₄, PTH₁₅₋₈₄, PTH₁₆₋₈₄, PTH₁₇₋₈₄, PTH₁₈₋₈₄, PTH₁₉₋₈₄, PTH₂₀₋₈₄, PTH₂₁₋₈₄, PTH₂₂₋₈₄, PTH₂₃₋₈₄, PTH₂₄₋₈₄, PTH₂₅₋₈₄, PTH₂₆₋₈₄, PTH₂₇₋₈₄, PTH₂₈₋₈₄, PTH₂₉₋₈₄, PTH₃₀₋₈₄, PTH₃₁₋₈₄, PTH₃₂₋₈₄, and PTH₃₃₋₈₄. In another specific embodiment, the PTH antagonist is a protein or a peptide, or a nucleic acid encoding said protein or peptide, selected from the group consisting of PTH₇₋₆₉, PTH₇₋₇₀, PTH₇₋₇₁, PTH₇₋₇₂, PTH₇₋₇₃, PTH₇₋₇₄, PTH₇₋₇₅, PTH₇₋₇₆, PTH₇₋₇₇, PTH₇₋₇₈, PTH₇₋₇₉, PTH₇₋₈₀, PTH₇₋₈₁, PTH₇₋₈₂, PTH₇₋₈₃ and PTH₇₋₈₄.

[0088] The PTH antagonist can have any suitable length provided that it maintains its antagonizing activity. For example, the PTH antagonist can have a length of 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45,

46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82 or 83 amino acid residues.

[0089] The PTH antagonist can further comprise an amino acid residue substitution or modification that enhances or does not decrease its antagonist activity, or an amino acid residue substitution or modification that stabilizes the PTH antagonist. For example, the PTH antagonist can further comprise the following amino acid residue substitution or modification: His₂₅, His₂₆, Leu₂₇, (U.S. Patent No. 5,382,658); Tyr₃₄, D-Trp₁₂, Nle_{8,18}, desamino(Nle_{8,18}), Lys₁₃ modified in the epsilon-amino acid group by N,N-diisobutyl or 3-phenylpropanoyl (U.S. Patent No. 5,093,233); Gly₁₂ substituted by D-Trp, L-Trp, L- or D- α - or β -naphthylalanine, or D- or L- α -MeTrp (U.S. Patent No. 4,968,669); the amino acid residue at positions 7, 11, 23, 24, 27, 28, or 31 being cyclohexylalanine, the amino acid residue at position 3, 16, 17, 18, 19, or 34 being α -aminoisobutyric acid, the amino acid residue at position 1 being α , β -diaminopropionic acid, the amino acid residue at position 27 being homoarginine, the amino acid residue at position 31 being norleucine (U.S. Patent No. 5,723,577); each of Arg₂₅, Lys₂₆, Lys₂₇ being substituted with Ala, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Met, Phe, Pro, Ser, Thr, Trp, Tyr or Val (U.S. Patent No. 5,317,010); and a combination thereof.

[0090] The PTH antagonist or a pharmaceutical composition comprising the same can be formulated in any suitable formats. For example, the PTH antagonist or a pharmaceutical composition comprising the same can be formulated in a solid or a liquid dosage form. In another example, the PTH antagonist or a pharmaceutical composition comprising the same can be formulated for oral, parenteral, intranasal, topical, or injectable administration, *e.g.*, intracavernous injection, subcutaneous injection, intravenous injection, intramuscular injection and intradermal injection.

[0091] If a nucleic acid encoding a PTH antagonist is used, the nucleic acid can be in any suitable forms, *e.g.*, DNA, RNA, PNA or a combination thereof. In a specific example, the nucleic acid can be comprised in a gene therapy vector, *e.g.*, an adenovirus associated vector, a retroviral vector, an adenovirus vector, and a lentivirus vector.

[0092] The PTH antagonist can be administered in any suitable regimen.

In one example, the PTH antagonist is administered as a bolus. In another example, the PTH antagonist is administered continuously. In still another example, the PTH antagonist is administered intermittently or is multiply

administered. In yet another example, the PTH antagonist is administered over a course of about 1, 2, 2-6, 6-12, or 12-24 hours. In yet another example, the PTH antagonist is administered over a course of about 1, 2, 2-5, 5-14, or 14-60 days. In yet another example, the PTH antagonist is administered over a course of about 1, 2, 2-6, 6-12, 12-24, 24-48, or more months. In yet another example, the PTH antagonist is administered intraperitoneally daily for about 2 months and then the administration is stopped for about 1 month and then resumed intraperitoneally daily for about 2 months.

[0093] In one specific embodiment, the human to be treated has undergone or is presently undergoing PTH agonist therapy, and the method is used to prevent, treat or delay excessive bone mineral, *e.g.*, calcium, loss caused by or associated with the PTH agonist therapy. Preferably, the method is used to prevent, treat or delay bone density decrease caused by or associated with said PTH agonist therapy. The PTH agonist used in the PTH agonist therapy can be any suitable PTH agonist, *e.g.*, the one having the PTH adenylate cyclase activating activity. Also preferably, the human to be treated has an abnormal PTH agonist and/or PTH antagonist level.

[0094] In another aspect, the present invention is directed to a method for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set forth in SEQ ID NO:2, dog PTH having an amino acid sequence set forth in SEQ ID NO:3, bovine PTH having an amino acid sequence set forth in SEQ ID NO:4, rat PTH having an amino acid

sequence set forth in SEQ ID NO:5, or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6, or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby said disease or disorder associated with excessive bone mineral, *e.g.*, calcium, loss is prevented, treated or delayed.

Methods for preventing, treating or delaying the effect of a PTH agonist

[0095] In another aspect, the present invention is directed to a method for preventing, treating or delaying the effect of a PTH agonist in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby the effect of said PTH agonist is prevented, treated or delayed.

[0096] The present method can be used for preventing, treating or delaying the effect of any PTH agonist, *e.g.*, the effect of a PTH agonist that has the PTH adenylate cyclase activating activity or for preventing, treating or delaying the effect of a human PTH agonist. Preferably, the PTH agonist to be

countered by the present method comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH agonist has the following characteristics: a) the N-terminal amino acid residue of said PTH agonist starts at position 1 of said PTH₁₋₈₄; and b) the C-terminal amino acid residue of said PTH agonist ends at any position spanning position 34 through position 84 of said PTH₁₋₈₄.

[0097] In a specific embodiment, the human to be treated has an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist. The abnormal comparative value between PTH agonist and PTH antagonist can be determined by any suitable formats, *e.g.*, as a ratio, a proportion or a subtraction value. In one example, the abnormal comparative value between PTH agonist and PTH antagonist is determined by determining and comparing at least two of the parameters selected from the group consisting of the level of the PTH agonist, the PTH antagonist and the total PTH level, *i.e.*, a sum of PTH agonist and PTH antagonist. In another example, the human to be treated has a PTH agonist/PTH antagonist ratio more than 2 and the method is used to bring the PTH agonist/PTH antagonist ratio within a range from about 1 to about 2.

[0098] The present method can be used for preventing, treating or delaying a disease or disorder associated with excessive PTH agonist effect. For example, the present method can be used for treating hypercalcemia, hyperparathyroidism, pseudohypoparathyroidism, a hypercalcemic crisis resulted from hyperparathyroidism, hyperparathyroidism caused by renal failure, a tumor producing a parathyroid hormone-like substance, an immune disease such as inflammation, an allergic response or hyperactive lymphocytes, or hypertension.

[0099] In one specific embodiment, the human to be treated is undergoing a calcium or vitamin D treatment, *e.g.*, for treating osteoporosis, and the method further comprises a step of monitoring the PTH antagonist level or a comparative value between PTH agonist and PTH antagonist to guide the administration of calcium or vitamin D in the patient.

[0100] In still another aspect, the present invention is directed to a method for preventing, treating or delaying the effect of a PTH agonist in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set forth in SEQ ID NO:2, dog PTH having an amino acid sequence set forth in SEQ ID NO:3, bovine PTH having an amino acid sequence set forth in SEQ ID NO:4, rat PTH having an amino acid sequence set forth in SEQ ID NO:5 or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6, or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby the effect of said PTH agonist is prevented, treated or delayed.

Methods for identifying risk of having osteoporosis or decreased bone density

[0101] In yet another aspect, the present invention is directed to a method for identifying a subject having or at risk of having osteoporosis or decreased bone density, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist as having or at risk of having osteoporosis or decreased bone density.

[0102] In a specific embodiment, the human to be treated has an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist. The abnormal comparative value between PTH agonist and PTH antagonist can be determined by any suitable formats, e.g., as a ratio, a proportion

or a subtraction value. In one example, the abnormal comparative value between PTH agonist and PTH antagonist is determined by determining and comparing at least two of the parameters selected from the group consisting of the level of the PTH agonist, the PTH antagonist and the total PTH level, *i.e.*, a sum of PTH agonist and PTH antagonist. In another example, the human to be treated has a PTH agonist/PTH antagonist ratio more than 2 and the method is used to bring the PTH agonist/PTH antagonist ratio within a range from about 1 to about 2.

Methods for identifying a subject in need of PTH antagonist treatment

[0103] In yet another aspect, the present invention is directed to a method for identifying a subject in need of parathyroid hormone (PTH) antagonist treatment, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist as in need of parathyroid hormone (PTH) antagonist treatment.

[0104] In a specific embodiment, the human to be treated has an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist. The abnormal comparative value between PTH agonist and PTH antagonist can be determined by any suitable formats, *e.g.*, as a ratio, a proportion or a subtraction value. In one example, the abnormal comparative value between PTH agonist and PTH antagonist is determined by determining and comparing at least two of the parameters selected from the group consisting of the level of the PTH agonist, the PTH antagonist and the total PTH level, *i.e.*, a sum of PTH agonist and PTH antagonist. In another example, the human to be treated has a PTH agonist/PTH antagonist ratio more than 2 and the method is used to bring the PTH agonist/PTH antagonist ratio within a range from about 1 to about 2, by using a PTH antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist to the identified subject.

[0105] Any suitable PTH antagonist, including the ones disclosed above, can be used in the present method. Preferably, the PTH antagonist comprises a

contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues.

D. The formulation, dosage and route of administration of PTH antagonist

[0106] The formulation, dosage and route of administration of PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, preferably in the form of pharmaceutical compositions, can be determined according to the methods known in the art (see *e.g.*, *Remington: The Science and Practice of Pharmacy*, Alfonso R. Gennaro (Editor) Mack Publishing Company, April 1997; *Therapeutic Peptides and Proteins: Formulation, Processing, and Delivery Systems*, Banga, 1999; and *Pharmaceutical Formulation Development of Peptides and Proteins*, Hovgaard and Frkjr (Ed.), Taylor & Francis, Inc., 2000; *Medical Applications of Liposomes*, Lasic and Papahadjopoulos (Ed.), Elsevier Science, 1998; *Textbook of Gene Therapy*, Jain, Hogrefe & Huber Publishers, 1998; *Adenoviruses: Basic Biology to Gene Therapy*, Vol. 15, Seth, Landes Bioscience, 1999; *Biopharmaceutical Drug Design and Development*, Wu-Pong and Rojanasakul (Ed.), Humana Press, 1999; *Therapeutic Angiogenesis: From Basic Science to the Clinic*, Vol. 28, Dole et al. (Ed.), Springer-Verlag New York, 1999). The PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, can be formulated for oral, rectal, topical, inhalational, buccal (*e.g.*, sublingual), parenteral (*e.g.*, subcutaneous, intramuscular, intradermal, or

intravenous), transdermal administration or any other suitable route of administration. The most suitable route in any given case will depend on the nature and severity of the condition being treated and on the nature of the particular PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, which is being used.

[0107] The PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, can be administered alone. Alternatively and preferably, the PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, is co-administered with a pharmaceutically acceptable carrier or excipient. Any suitable pharmaceutically acceptable carrier or excipient can be used in the present method (*See e.g., Remington: The Science and Practice of Pharmacy*, Alfonso R. Gennaro (Editor) Mack Publishing Company, April 1997).

[0108] The present method can be used alone. Alternatively, the present method can be used in combination with other agent suitable for preventing, treating or delaying a disease or disorder associated with excessive bone mineral, e.g., calcium, loss. Such other agent can be used before, with or after the administration of PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist. For example, the PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, can be co-administered with such other agent.

[0109] The nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, can be used in the form of naked DNA, complexed DNA, cDNA, plasmid DNA, RNA or other mixtures thereof as components of the

gene delivery system. In another embodiment, the nucleic acid encoding a PTH antagonist protein, or a functional fragment thereof, is included in a viral vector. Any viral vectors that are suitable for gene therapy can be used in the combination. For example, an adenovirus vector (U.S. Patent No. 5,869,305), a simian virus vector (U.S. Patent No. 5,962,274), a conditionally replicating human immunodeficiency viral vector (U.S. Patent No. 5,888,767), retrovirus, SV40, Herpes simplex viral amplicon vectors and Vaccinia virus vectors can be used. In addition, the genes can be delivered in a non-viral vector system such as a liposome wherein the lipid protects the DNA or other biomaterials from oxidation during the coagulation.

[0110] According to the present invention, the PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, alone or in combination with other agents, carriers or excipients, may be formulated for any suitable administration route, such as intracavernous injection, subcutaneous injection, intravenous injection, intramuscular injection, intradermal injection, oral or topical administration. The method may employ formulations for injectable administration in unit dosage form, in ampoules or in multidose containers, with an added preservative. The formulations may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, sterile pyrogen-free water or other solvents, before use. Topical administration in the present invention may employ the use of a foam, gel, cream, ointment, transdermal patch, or paste.

[0111] Pharmaceutically acceptable compositions and methods for their administration that may be employed for use in this invention include, but are not limited to those described in U.S. Patent Nos. 5,736,154; 6,197,801 B1; 5,741,511; 5,886,039; 5,941,868; 6,258,374 B1; and 5,686,102.

[0112] The magnitude of a therapeutic dose in the treatment or prevention will vary with the severity of the condition to be treated and the route of administration. The dose, and perhaps dose frequency, will also vary according to age, body weight, condition and response of the individual patient.

5 [0113] It should be noted that the attending physician would know how to and when to terminate, interrupt or adjust therapy to lower dosage due to toxicity, or adverse effects. Conversely, the physician would also know how to and when to adjust treatment to higher levels if the clinical response is not adequate (precluding toxic side effects).

10 [0114] Any suitable route of administration may be used. Dosage forms include tablets, troches, cachet, dispersions, suspensions, solutions, capsules, patches, and the like. See, Remington's Pharmaceutical Sciences.

[0115] In practical use, the PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a
15 functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, alone or in combination with other agents, may be combined as the active in intimate admixture with a pharmaceutical carrier or excipient, such as beta-cyclodextrin and 2-hydroxy-propyl-beta-cyclodextrin, according to conventional pharmaceutical compounding
20 techniques. The carrier may take a wide form of preparation desired for administration, topical or parenteral. In preparing compositions for parenteral dosage form, such as intravenous injection or infusion, similar pharmaceutical media may be employed, water, glycols, oils, buffers, sugar, preservatives, liposomes, and the like known to those of skill in the art. Examples of such
25 parenteral compositions include, but are not limited to dextrose 5% w/v, normal saline or other solutions. The total dose of the PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, alone or in combination with other
30 agents to be administered may be administered in a vial of intravenous fluid,

ranging from about 1 ml to 2000 ml. The volume of dilution fluid will vary according to the total dose administered.

[0116] The invention also provides for kits for carrying out the therapeutic regimens of the invention. Such kits comprise in one or more containers therapeutically effective amounts of the PTH antagonist protein, or a functional fragment thereof, or a nucleic acid encoding an PTH antagonist protein, or a functional fragment thereof, or an agent that enhances production and/or antagonizing function of said PTH antagonist, alone or in combination with other agents, in pharmaceutically acceptable form. Preferred pharmaceutical forms would be in combination with sterile saline, dextrose solution, or buffered solution, or other pharmaceutically acceptable sterile fluid. Alternatively, the composition may be lyophilized or dessicated; in this instance, the kit optionally further comprises in a container a pharmaceutically acceptable solution, preferably sterile, to reconstitute the complex to form a solution for injection purposes. Exemplary pharmaceutically acceptable solutions are saline and dextrose solution.

[0117] In another embodiment, a kit of the invention further comprises a needle or syringe, preferably packaged in sterile form, for injecting the composition, and/or a packaged alcohol pad. Instructions are optionally included for administration of composition by a physician or by the patient.

[0118] In specific embodiments, formulations disclosed in the following U.S. Patent Nos. can be used. U.S. Patent No. 5,496,801 discloses a PTH formulation using mannitol as excipient and citrate as buffering agent, and are incorporated in vials as a freeze-dried powder for reconstitution to treat osteoporosis. U.S. Patent No. 5,607,915 discloses systemic delivery of parathyroid hormone to a mammalian host by inhalation through the mouth of a dispersion of an N-terminal fragment of PTH. It has been found that such respiratory delivery of the PTH fragment provides a pulsatile concentration profile of the PTH in the host's serum. PTH fragment compositions include dry powder formulations having the PTH present in a dry bulking powder, liquid solutions or suspensions suitable for nebulization, and aerosol propellants suitable for use in a

metered dose inhaler. U.S. Patent No. 5,563,122 discloses a safe, stabilized lyophilized preparation comprising PTH as an active ingredient and effective amounts of sugar and sodium chloride to provide a stable preparation of parathyroid hormone. The sugar can be a monosaccharide, *e.g.*, mannitol, glucose, sorbitol, inositol, and or a disaccharide, *e.g.*, sucrose, maltose, lactose or trehalose.

E. Methods for monitoring treatment for osteoporosis or decreased bone density

10 [0119] In yet another aspect, the present invention is directed to a method for monitoring a subject undergoing treatment for osteoporosis or decreased bone density, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist.

15 [0120] The abnormal comparative value between PTH agonist and PTH antagonist can be determined by any suitable formats, *e.g.*, as a ratio, a proportion or a subtraction value. In one example, the abnormal comparative value between PTH agonist and PTH antagonist is determined by determining and comparing at least two of the parameters selected from the group consisting of the level of the PTH agonist, the PTH antagonist and the total PTH level, *i.e.*, a sum of PTH agonist and PTH antagonist. In another example, the human to be treated has a PTH agonist/PTH antagonist ratio more than 2. In still another example, the subject, *e.g.*, a human, has PTH agonist-PTH antagonist value that equals or is more than 50 pg/ml. In yet another example, the subject, *e.g.*, a human, has a PTH agonist level that is more than PTH antagonist level. The present method can further comprise a step to bring the PTH agonist/PTH antagonist ratio within a range from about 1 to about 2, by using a PTH antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist to the identified subject.

[0121] In a specific embodiment, the subject, e.g., a human, is undergoing a calcium, bisphosphanate or vitamin D treatment. This way, the present method can be used to guide the calcium, bisphosphanate or vitamin D treatment in order for those agents to be useful in stimulating the correct amount of PTH antagonist as an osteoclast inhibitor for the treatment of osteoporosis and high bone turnover. The new discovery here is that calcium and vitamin D cause a preferential stimulation of PTH antagonist which acts directly on the bone to lower bone turnover by inhibiting the osteoclasts (Divietti et al., *Endocrinology*, 143:171-176 (2002); and Divietti et al., *J. Bone Miner. Res.*, 16:Suppl 1, S307 (2001)).

Accordingly, the present method can be used to monitor the effectiveness of the calcium, bisphosphanate or vitamin D treatment for osteoporosis and high bone turnover by measuring both the absolute increase in PTH antagonist and the increase in PTH antagonist relative to PTH agonist.

[0122] Any suitable PTH antagonist, including the ones disclosed above, can be used in the present method. Preferably, the PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues.

F. Methods for identifying an agent for preventing, treating or delaying osteoporosis

[0123] In yet another aspect, the present invention is directed to a method for identifying an agent suitable for preventing, treating or delaying osteoporosis, which method comprises: a) measuring PTH antagonist activity in the presence and absence of a test substance; and b) identifying a substance that enhances said

PTH antagonist activity as an agent suitable for preventing, treating or delaying osteoporosis.

[0124] Any suitable PTH antagonist, including the ones disclosed above, can be used in the present method. Preferably, the PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues.

[0125] In comparing the activity of a PTH antagonist in the presence and absence of a test substance to assess whether the test substance is a potentiator of the PTH antagonist, it is unnecessary to assay the activity in parallel, although such parallel measurement is preferred. It is possible to measure the activity of the PTH antagonist at one time point and compare the measured activity to a historical value of the activity of the PTH antagonist. For instance, one can measure the activity of the PTH antagonist in the presence of a test substance and compare with historical value of the activity of the PTH antagonist measured previously in the absence of the test substance, and *vice versa*. This can be accomplished, for example, by providing the activity of the PTH antagonist on an insert or pamphlet provided with a kit for conducting the assay.

[0126] A variety of formats and detection protocols are known for performing screening assays. Any such formats and protocols may be adapted for identifying potentiator of PTH antagonist activities. The following includes a discussion of exemplary protocols.

High throughput screening assays

[00100] Although the above-described assay can be conducted where a single PTH antagonist is screened against, and/or a single test substance is

screened for in one assay, the assay is preferably conducted in a high throughput screening mode, *i.e.*, a plurality of the PTH antagonists are screened against and/or a plurality of the test substances are screened for simultaneously (*See generally*, High Throughput Screening : The Discovery of Bioactive Substances (Devlin, Ed.) Marcel Dekker, 1997; Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997); and Silverman et al., *Curr. Opin. Chem. Biol.*, 2(3):397-403 (1998)). For example, the assay can be conducted in a multi-well (*e.g.*, 24-, 48-, 96-, or 384-well), chip or array format. High-throughput screening (HTS) is the process of testing a large number of diverse chemical structures against disease targets to identify "hits" (Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997)). Current state-of-the-art HTS operations are highly automated and computerized to handle sample preparation, assay procedures and the subsequent processing of large volumes of data. Detection technologies employed in high-throughput screens depend on the type of biochemical pathway being investigated (Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997)). These methods include, radiochemical methods, such as the scintillation-proximity assays (SPA), which can be adapted to a variety of enzyme assays (Lerner et al., *J. Biomol. Screening*, 1:135-143 (1996); Baker et al., *Anal. Biochem.*, 239:20-24 (1996); Baum et al., *Anal. Biochem.*, 237:129-134 (1996); and Sullivan et al., *J. Biomol. Screening*, 2:1923 (1997)) and protein-protein interaction assays (Braunwalder et al., *J. Biomol. Screening*, 1:23-26 (1996); Sonatore et al., *Anal. Biochem.*, 240:289-297 (1996); and Chen et al., *J. Biol. Chem.*, 271:25308-25315 (1996)), and nonisotopic detection methods, including but are not limited to, colorimetric and luminescence detection methods, resonance energy transfer (RET) methods, time-resolved fluorescence (HTRF) methods, cell-based fluorescence assays, such as fluorescence resonance energy transfer (FRET) procedures (*see, e.g.*, Gonzalez et al., *Biophys. J.*, 69:1272-1280 (1995)), fluorescence polarization or anisotropy methods (*see, e.g.*, Jameson et al., *Methods Enzymol.*, 246:283-300 (1995); Jolley, *J. Biomol. Screening*, 1:33-38 (1996); Lynch et al., *Anal. Biochem.*, 247:77-82 (1997)), fluorescence correlation spectroscopy (FCS) and other such methods.

Test Substances

[0127] Test compounds, including small molecules and libraries and collections thereof can be screened in the above-described assays and assays described below to identify compounds that potentiate the activity a PTH antagonist. Rational drug design methodologies that rely on computational chemistry may be used to screen and identify candidate compounds. The compounds identified by the screening methods include any compounds and collections of compounds available, known or that can be prepared.

[0128] Compounds can be selected for their potency and selectivity of potentiation of a PTH antagonist. The assay is performed in the absence of test compound, and in the presence of increasing concentrations of the test compound. The concentration of test compound at which 50% of the PTH antagonist activity is potentiated by the test compound is the IC₅₀ value (Potentiation Concentration) or EC₅₀ (Effective Concentration) value for that compound. Within a series or group of test compounds, those having lower IC₅₀ or EC₅₀ values are considered more potent potentiators of the PTH antagonist than those compounds having higher IC₅₀ or EC₅₀ values. The IC₅₀ measurement is often used for more simplistic assays, whereas the EC₅₀ is often used for more complicated assays, such as those employing cells.

[0129] Preferred compounds according to this aspect have an IC₅₀ value of 100 nM or less as measured in an *in vitro* assay for potentiation of a PTH antagonist. Especially preferred compounds have an IC₅₀ value of less than 100 nM.

[0130] The test compounds also are evaluated for selectivity toward a PTH antagonist. As described herein, and as generally known, a test compound is assayed for its potency toward a panel of target PTH antagonists and other PTH related moieties and an IC₅₀ value or EC₅₀ value is determined for each test compound in each assay system. A compound that demonstrates a low IC₅₀ value or EC₅₀ value for the target PTH antagonist, and a higher IC₅₀ value or EC₅₀ value for the other PTH related moieties within the test panel, is considered to be selective toward the target PTH antagonist. Generally, a compound is

deemed selective if its IC50 value or EC50 value in the target PTH antagonist assay is at least one order of magnitude less than the next smallest IC50 value or EC50 value measured in the selectivity panel of PTH antagonists.

[0131] Compounds are also evaluated for their activity *in vivo*. The type of assay chosen for evaluation of test compounds will depend on the pathological condition to be treated or prevented by use of the compound, as well as the route of administration to be evaluated for the test compound.

G. Methods and kits for preventing, treating or delaying bone metastasis

[0132] In yet another aspect, the present invention is directed to a method for preventing, treating or delaying bone metastasis in a human, which method comprises administering to a human, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics: a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄; b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and c) said PTH antagonist has a minimal length of three amino acid residues, whereby said bone metastasis is prevented, treated or delayed.

[0133] The present methods can be used to prevent, treat or delay any bone metastasis, *e.g.*, the bone metastasis associated with or caused by breast cancer, prostate cancer or multiple myeloma. For example, the present methods can be used to prevent, treat or delay morbidity or mortality associated with the bone metastasis. In another example, the present methods can be used to prevent, treat or delay pain, pathological fractures, hypercalcemia or spinal cord compression associated with the bone metastasis.

[0134] Any suitable PTH antagonists or agents that enhances production and/or antagonizing function of the PTH antagonists, including the ones described in the present application, can be used in the present methods. The PTH antagonists or agents that enhances production and/or antagonizing function of the PTH antagonists can be administered via any suitable route. For example, the PTH antagonists or an agents that enhances production and/or antagonizing function of the PTH antagonists can be administered orally or intravenously.

[0135] The PTH antagonists or agents that enhances production and/or antagonizing function of the PTH antagonists can be used alone in the present methods. Alternatively, the PTH antagonists or agents that enhances production and/or antagonizing function of the PTH antagonists can be used in combination with a hormone therapy or a chemotherapy.

[0136] In yet another aspect, the present invention is directed to a kit, which kit comprises an above-described parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of the PTH antagonist in a container and an instruction for using the PTH antagonist or agent in preventing, treating or delaying bone metastasis.

H. Exemplary embodiments

[0137] There are a number of closely analogous, species dependent forms of PTH. The amino acid sequence of hPTH or cyclase activating parathyroid hormone (CAP) is shown in FIGURE 1. However, for rat PTH, bovine PTH, or porcine PTH, for example, one finds the substitutions at some of the amino acids in the hPTH sequence. For the purposes of the present embodiment, one can use interchangeably truncated forms of these PTH's, although it is preferred to use a PTH having a sequence matching the species in which the PTH antagonist is used.

[0138] Preferred PTH antagonists of the present embodiment have an amino acid sequence from between PTH₂₋₈₄ and PTH₃₄₋₈₄ or a conservatively substituted variant thereof exhibiting PTH antagonist activity, with the most preferred form being from between PTH₃₋₈₄ and PTH₂₈₋₃₄.

[0139] In order to make the present compositions, one can use any conventionally known method. For example, one can use recombinant DNA methods to produce the desired compound. Alternatively, one can use an automated peptide synthesizer, such as Model 431 made by Applied Biosystems, Inc. (Foster City, California, U.S.A.) Fmoc (9-fluoronylmethoxycarbonyl) can be used as the alpha-amino protecting group. All amino acids and solvents are available from Applied Biosystems and are of synthesis grade. Following synthesis, the peptide is cleaved from the resin, and side chains are de-blocked, using a cleavage cocktail containing 6.67% phenol, 4.4% (v/v) thioanisole and 8.8% ethanedithiol in trifluoroacetic acid (TFA). The cleaved peptide is precipitated and washed several times in cold diethyl ether. It is then dissolved in water and lyophilized. The crude peptide is subjected to amino acid analysis (Waters PICO-TAG System, Boston, Massachusetts, U.S.A.) and reversed-phase HPLC using a VYDAC (TM) C8 column with 0.1% TFA in water and 99.9% acetonitrile in 0.1% TFA as the mobile buffers. The presence of a single major peak along with the appropriate amino acid composition is taken as evidence that the peptide is suitable for further use.

[0140] The exemplary PTH antagonist peptides exhibit both oral and parenteral activity and can be formulated in solid or liquid dosage forms for oral, parenteral, intranasal, topical, or injectable administration using known carriers, excipients, or the like. The exact amount of exemplary PTH antagonist used can vary depending upon the degree of antagonist property desired, the route of administration, or the duration of the treatment, as is known to the art.

[0141] The exemplary PTH antagonists have the ability to reduce the increase in serum calcium normally caused by PTH or a PTH agonist analog. These antagonists also possess an ability similar to a cyclase inhibiting parathyroid hormone peptide (CIP) to treat osteoporosis due to the CAP rebound effect that comes with CIP administration. The CAP rebound effect is believed to be the body's response to an administration of CIP. This response occurs when the parathyroid gland cells secrete CAP in an effort to return the CAP/CIP ratio to homeostasis with the pre-CIP administration levels. PTH antagonist

administration is not accompanied by hypercalcemia and osteosarcoma, as with direct CAP administration. FIGURE 5 is a graph demonstrating such a hypercalcemic prevention property. Twenty-five rats were used in a demonstration of the effect of the exemplary PTH antagonists. All of the rats had their parathyroid glands removed. Five rats received an *i.v.* injection of a saline control. The serum calcium of the control rats was measured and on average was lowered over time by about 0.18 mg/dl by virtue of a parathyroidectomy. Nine rats received an *i.v.* injection (10 µg/kg) of hPTH obtained from Bachem, AG of Bubendorf, Switzerland. The serum calcium of the hPTH rats was measured and on average was raised over time by about 0.65 mg/dl. Five rats received an equimolar *i.v.* injection of a PTH₇₋₈₄ (an exemplary PTH antagonist) also obtained from Bachem, AG of Bubendorf, Switzerland. The serum calcium of the PTH antagonist rats was measured and on average was lowered over time by about 0.30 mg/dl. Finally, six rats received an *i.v.* injection comprised of hPTH (10 µg/kg) and an equimolar amount of PTH antagonist PTH₇₋₈₄. The serum calcium of the hPTH/PTH antagonist rats was measured and on average remained substantially the same over time, raising only about 0.03 mg/dl. Thus, the exemplary composition was able to prevent the substantial serum calcium increase normally associated with an administration of hPTH to rats having hypoparathyroidism, and quite unexpectedly, is much more potent in its antagonist property than the previously reported antagonist PTH₃₋₃₄.

I. Examples

METHODS

25 Characterization of the new "Whole" PTH IRMA Assay

[0142] This new Whole PTH two-site assay (Scantibodies Laboratories, Santee, CA, USA) first employees and antibody that recognizes the 39-84 region of the PTH molecule. This antibody, produced in a goat and affinity purified, is present in relative excess and is immobilized onto polystyrene-coated tubes. The

- second antibody, also developed in a goat, was also affinity purified and recognizes only the first six amino acids (1 to 6; Ser-Val-Ser-Glu-Ile-Gln) of the human PTH molecule (Fig. 1). This anti-hPTH assay uses synthetic human PTH 1-84 as the standard, with a limit of detection of approximately 1 to 2 pg/mL.
- 5 Normal values range from 5 to 35 pg/mL. The interassay and intra-assay coefficients of variation were found to be between 2 and 7% and recovery was from 96 to 106%. The Whole PTH assay was compared with the Intact PTH assay purchases from the Nichols Institute (I-Nichols, San Jan Capistrano, CA, USA). Synthetic human PTH 1-84 and 7-84 were purchased from Bachem
- 10 (Torrance, CA USA). To assess circulating levels of hPTH 1-84 and non-(1-84) PTH, heparinized blood samples were obtained before dialysis from 28 patients who had been maintained on chronic hemodialysis for 1.2 to 7.5 years and from 14 renal transplant patients (1 to 6 years).

Studies *in vitro*

- 15 [0143] *Osteoblastic cell line.* To compare the biological effects of the two peptides (hPTH 1-84 and 7-84), intracellular cAMP production was measured in the rate osteosarcoma cell line ROS/17.2, which has an osteoblastic phenotype and is known to increase cAMP production in response to PTH. Cells were cultured in Ham's F12 media containing 10% fetal bovine serum. Cells were
- 20 plated out in 12-well plates at a density of 30,000 cells per well and grown to confluence. Cells were washed three times with KHMS buffer at 37°C (KCl 4.0 mmol/L, CaCl₂ 1.25 mmol/L, MgSO₄ 1.25 mmol/L, KH₂PO₄ 1.2 mmol/L, HEPES 10 mmol/L, NaCl 100 mmol/L, NaHCO₃ 37 mmol/L, and glucose 10 mmol/L, pH 7.5). cAMP production was measured using 500 µL of
- 25 KHMS buffer (37°) containing isobutyl-1 methylxanthine (IBMX) 1.0 mmol/L and various concentrations (10⁻¹¹ to 10⁻⁸ mol/L) of hPTH 1-84 or hPTH 7-84. After a five-minute incubation, 100 µL of 1.8 mol/L perchloric acid were added. After an additional five-minute incubation at room temperature, 100 µL of 3 mol/KHCO₃ were added to neutralize the acid. Samples were centrifuged at 3000 rpm for 15
- 30 minutes, and the supernatants were assayed for cAMP [26].

[0144] *Analysis of PTH in human parathyroid glands.* Human parathyroid glands were placed in ice-cold phosphate-buffered saline and processed within 30 minutes of parathyroidectomy. Aliquots of parathyroid tissue were dissected, weighed, and homogenized in 500 μ L of a buffer containing 100 mmol/L Tris-HCl, pH 7.5, 100 mmol/L NaCl, 1 mol/L DL-dithiothreitol, and a complete TM protease inhibitor cocktail (Boehringer-Mannheim, Mannheim, Germany). Homogenates were sonicated three times for 30 seconds each at 0°C and centrifuged at 10,000 X g for 15 minutes. Supernatants were kept at -70°C until measurements of 1-84 PTH, non-(1-84) PTH, and total protein were performed.

10 Studies in vivo

[0145] *Calcemic response.* Normal female Sprague-Dawley rats weighing 225 to 250 g (Harlan, Indianapolis, IN, USA) were parathyroidectomized (PTX) and fed a 0.02% calcium diet. Rats with a plasma calcium below 7.0 mg/dL after overnight fasting were included in the study. A 20 μ g dose of hPTH 1-84 or 7-84 was given intraperitoneally to PTX rats in four doses of 5 μ g each at 30-minute intervals (0, 30, 60, and 90 minutes). For control studies, the rats received vehicle (saline solution) alone. Blood was drawn via the tail at 0, 60, 90, and 120 minutes. For competition experiments, rats received an injection of hPTH 7-84 10 minutes prior to each injection of hPTH 1-84. The molar ratio of hPTH 7-84:hPTH 1-84 was 1:1.

[0146] *Phosphaturic response.* Normal female Sprague-Dawley rats weighing 225 to 250 g were prepared for clearance studies under light anesthesia. Polyethylene catheters (PE50) were placed in the femoral artery for the collection of blood and measurement of blood pressure (Blood Pressure Analyzer; Micro-Medic, Inc., Louisville, KY, USA), in the femoral vein for infusion and in the bladder for the collection of urine. Rats were placed in Plexiglas® holders and allowed to recover from the effect of the anesthetic for one hour. A priming dose (0.6 mL) of chemical inulin in saline was administered over a period of three minutes to achieve a plasma inulin level between 50 and 100 mg/mL. A solution of saline containing inulin to maintain this level and calcium gluconate to deliver

0.5 mg. calcium was infused at the rate of 0.03 mL/min. After equilibration, a total of four 30-minute urine collections was obtained.

[0147] To assess the effect of hPTH 1-84 on phosphate excretion, urine was collected during two control periods, after which rats received a priming bolus of 1.8 μ g of hPTH 1-84 followed by a sustained infusion that delivered a total of 8.2 μ g of I-PTH. After an equilibration period of 20 minutes, two 30-minute urine collections were obtained. In competition experiments, hPTH 7-84 was given five minutes prior to hPTH 1-84 at a molar ratio of 4:1.

[0148] Blood samples and blood pressure measurements were recorded at the beginning and end of the baseline period, at the beginning of the PTH infusion period, and at the end of the study. The concentration of inulin in plasma and urine was determined by the method of Führ, Kaczmarczyk, and Kruttgen, *Klin Wochenschr*, 33:729-730 (1955). The estimation of the glomerular filtration rate (GFR) by inulin clearance and the calculation of the fractional urinary excretion rate of phosphorus (FE_{P04}) were performed in the standard fashion. Blood samples were centrifuged, and plasma phosphorus and calcium concentrations were measured.

Serum chemistries

[0149] Total plasma calcium levels were determined using an atomic absorption spectrophotometer (model 1100B; Perkin Elmer, Norwalk, CT, USA). Plasma phosphorus levels were determined using an autoanalyzer (COBAS MIRA Plus; Roche, Newark, NJ, USA).

Statistical analysis

[0150] Results are expressed as mean \pm SEM. *N* indicates the number of samples. The paired *t*-test was employed to examine statistical significance, unless otherwise indicated in the text.

RESULTS

Specificity of IRMA assays for hPTH 1-84

[0151] Initial studies compared the ability of the Nichols Intact (I-Nichols) PTH assay and the new Whole PTH assay to discriminate between the hPTH 1-84 and hPTH 7-84 molecules. Figure 2 shows that the Nichols "intact" PTH assay did not discriminate between human PTH 1-84 and 7-84. However, as depicted in Figure 3, studies performed using the Whole PTH assay show that hPTH 1-84 was detected with a high degree of sensitivity, whereas hPTH 7-84 was undetectable, even at a concentration as high as 10,000 pg/mL.

Studies *in vitro*

[0152] The results of cAMP production by ROS/17.2 cells exposed to hPTH 1-84 or hPTH 7-84 are shown in Figure 4. Unlike hPTH 7-84, hPTH 1-84 increased cAMP production in a dose-dependent manner. hPTH 1-84 (10^{-8} mol/L) increased intracellular cAMP from 18.1 ± 1.25 to 738 ± 4.13 nmol/well. On the other hand, the same concentration of hPTH 7-84 had no effect on cAMP ($N = 6$).

Studies *in vivo* in rats

[0153] We next examined the hPTH 7-84 fragment as a potential competitive inhibitor of hPTH 1-84 in bone by measuring changes in serum calcium in PTX rats. Figure 5 shows that the administration of hPTH 1-84 to PTX rats fed a 0.02% calcium diet increased plasma calcium by 0.65 ± 0.10 mg/dL ($N=9$, $P < 0.001$, ANOVA). With the administration of vehicle alone, plasma calcium changed slightly in accordance with PTX (-0.17 ± 0.10 mg/dL, $N = 5$). A slight but significant decrease was observed in the rats receiving hPTH 7-84 (0.30 ± 0.08 mg/dL, $N = 5$, $P < 0.05$). When both peptides were given together in a 1:1 molar ratio, the calcemic response induced by the administration of hPTH 1-84 alone decreased by 94% ($N = 6$, $P < 0.001$, ANOVA). Thus in this model, hPTH 7-84 significantly inhibits hPTH 1-84 induction of bone calcium mobilization.

[0154] The phosphaturic effects of these two peptides were then evaluated (Fig. 6). The GFR did not change in rats infused with hPTH 1-84 (1.8 ± 0.3 vs. 1.8 ± 0.1 mL/min), whereas fractional excretion of phosphate (FE_{P04}) increased from 11.9 ± 2.4 to $27.7 \pm 2.4\%$ ($N = 10$, $P < 0.001$). When hPTH 7-84 was given simultaneously with hPTH 1-84, the GFR increased from 2.1 ± 0.1 to 2.6 ± 0.2 mL/min ($N = 8$, $P < 0.05$). However, despite this increase in GFR, the increase in FE_{P04} induced by treatment with hPTH 1-84 was significantly decreased by 50.2% ($P < 0.01$), by virtue of the coadministration of hPTH 7-84.

Studies in humans

[0155] Figure 7 shows that the values for plasma PTH were higher in all 28 patients on chronic dialysis when measured with the I-Nichols assay compared with the Whole assay. The median PTH values were 523 versus 344 pg/ml ($P < 0.001$), respectively. A regression analysis of these data is shown in Figure 8.

[0156] The association between plasma levels of non-(1-84) PTH, "likely" hPTH 7-84, and plasma calcium and phosphorus was next examined in 20 patients maintained on chronic dialysis (Fig. 9). There was a positive correlation between the percentage of non-(1-84) PTH and serum calcium ($P < 0.002$), but no correlation with plasma phosphorus (data not shown). These studies were performed only in those patients in whom there were values for calcium, phosphorous, and PTH from the same blood sample [20].

[0157] In a group of 14 renal transplant patients the percentage of non-(1-84) PTH was found to be $44.1 \pm 3.1\%$ of the total PTH, as measured by the I-Nichols assay and the Whole PTH assay (Fig. 10). The absolute PTH value with the I-Nichols assay was 132.9 ± 39.9 compared with 79.8 ± 24.8 pg/mL ($P < 0.005$) with the Whole PTH assay.

[0158] Finally, we examined whether intracellular cleavage of the hPTH 1-84 molecule occurs in the parathyroid gland, thus producing the non (1-84) PTH fragment. Surgically excised parathyroid glands from six uremic patients maintained on chronic dialysis were studied. Figure 11 shows that non (1-84) PTH fragments exist in the cell lysates from these parathyroid glands and

represent $41.8 \pm 3.2\%$ ($P < 0.05$) of the total intracellular PTH measured by the "intact" PTH assay (that is, 1-84 PTH and most likely 7-84 PTH).

[0159] The above examples are included for illustrative purposes only and are not intended to limit the scope of the invention. Many variations to those
5 described above are possible. Since modifications and variations to the examples described above will be apparent to those of skill in this art, it is intended that this invention be limited only by the scope of the appended claims.

CLAIMS

1. A pharmaceutical composition, which pharmaceutical composition comprises an effective amount of a parathyroid hormone (PTH) antagonist and a pharmaceutically acceptable carrier or excipient, wherein said PTH antagonist
 - 5 comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics:
 - a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
 - 10 b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
 - c) said PTH antagonist has a minimal length of three amino acid residues.
2. The pharmaceutical composition of claim 1, wherein the N-terminal
 - 15 amino acid residue of the PTH antagonist starts at position 2 of the PTH₁₋₈₄.
3. The pharmaceutical composition of claim 1, wherein the C-terminal
 - amino acid residue of the PTH antagonist ends at position 84 of the PTH₁₋₈₄.
- 20 4. The pharmaceutical composition of claim 1, wherein the PTH antagonist is a protein or a peptide, or a nucleic acid encoding said protein or peptide, selected from the group consisting of PTH₂₋₈₄, PTH₃₋₈₄, PTH₄₋₈₄, PTH₅₋₈₄, PTH₆₋₈₄, PTH₇₋₈₄, PTH₈₋₈₄, PTH₉₋₈₄, PTH₁₀₋₈₄, PTH₁₁₋₈₄, PTH₁₂₋₈₄, PTH₁₃₋₈₄, PTH₁₄₋₈₄, PTH₁₅₋₈₄, PTH₁₆₋₈₄, PTH₁₇₋₈₄, PTH₁₈₋₈₄, PTH₁₉₋₈₄, PTH₂₀₋₈₄, PTH₂₁₋₈₄, PTH₂₂₋₈₄,
 - 25 PTH₂₃₋₈₄, PTH₂₄₋₈₄, PTH₂₅₋₈₄, PTH₂₆₋₈₄, PTH₂₇₋₈₄, PTH₂₈₋₈₄, PTH₂₉₋₈₄, PTH₃₀₋₈₄, PTH₃₁₋₈₄, PTH₃₂₋₈₄, and PTH₃₃₋₈₄.
5. The pharmaceutical composition of claim 1, wherein the PTH antagonist is a protein or a peptide, or a nucleic acid encoding said protein or
 - 30 peptide, selected from the group consisting of PTH₇₋₆₉, PTH₇₋₇₀, PTH₇₋₇₁, PTH₇₋₇₂,

PTH₇₋₇₃, PTH₇₋₇₄, PTH₇₋₇₅, PTH₇₋₇₆, PTH₇₋₇₇, PTH₇₋₇₈, PTH₇₋₇₉, PTH₇₋₈₀, PTH₇₋₈₁,
PTH₇₋₈₂, PTH₇₋₈₃ and PTH₇₋₈₄.

6. The pharmaceutical composition of claim 1, wherein the PTH
5 antagonist has a length of 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19,
20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41,
42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63,
64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82 or 83
amino acid residues.

10

7. The pharmaceutical composition of claim 1, wherein the PTH
antagonist further comprises an amino acid residue substitution or modification
that enhances or does not decrease its antagonist activity, or an amino acid residue
substitution or modification that stabilizes the PTH antagonist.

15

8. The pharmaceutical composition of claim 7, wherein the amino acid
residue substitution or modification is selected from the group consisting of His₂₅,
His₂₆, Leu₂₇, Tyr₃₄, D-Trp₁₂, Nle_{8,18}, desamino(Nle_{8,18}), Lys₁₃ modified in the
epsilon-amino acid group by N,N-diisobutyl or 3-phenylpropanoyl, Gly₁₂
20 substituted by D-Trp, L-Trp, L- or D- α - or β -naphthylalanine, or D- or L- α -
MeTrp, the amino acid residue at positions 7, 11, 23, 24, 27, 28, or 31 being
cyclohexylalanine, the amino acid residue at position 3, 16, 17, 18, 19, or 34 being
 α -aminoisobutyric acid, the amino acid residue at position 1 being α , β -
diaminopropionic acid, the amino acid residue at position 27 being homoarginine,
25 the amino acid residue at position 31 being norleucine, each of Arg₂₅, Lys₂₆, Lys₂₇
being substituted with Ala, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Met, Phe,
Pro, Ser, Thr, Trp, Tyr or Val, and a combination thereof.

9. The pharmaceutical composition of claim 1, which is formulated in a
30 solid or a liquid dosage form.

10. The pharmaceutical composition of claim 1, which is formulated for oral, parenteral, intranasal, topical, or injectable administration.

11. The pharmaceutical composition of claim 10, wherein the injectable
5 administration is selected from the group consisting of intracavernous injection, subcutaneous injection, intravenous injection, intramuscular injection and intradermal injection.

12. The pharmaceutical composition of claim 1, wherein the nucleic acid
10 is a DNA.

13. The pharmaceutical composition of claim 1, wherein the nucleic acid is an RNA.

14. The pharmaceutical composition of claim 1, wherein the nucleic acid
15 is comprised in a gene therapy vector.

15. The pharmaceutical composition of claim 14, wherein the gene therapy
20 vector is selected from the group consisting of an adenovirus associated vector, a retroviral vector, an adenovirus vector, and a lentivirus vector.

16. A kit, which kit comprises a pharmaceutical composition of claim 1 in
a container and an instruction for using the pharmaceutical composition in
preventing, treating or delaying a disease or disorder associated with excessive
25 bone mineral loss or for balancing the effect of excess PTH agonist.

17. The kit of claim 16, wherein the disease or disorder associated with
excessive bone mineral loss is selected from the group consisting of
hyperparathyroidism, renal osteodystrophy, osteoporosis, parathyroid cancer,
30 hypercalcemia, an immune disease and hypertension.

18. A combination, which combination comprises an effective amount of a parathyroid hormone (PTH) antagonist and an effective amount of an agent suitable for preventing, treating or delaying a disease or disorder associated with excessive bone mineral loss, wherein said PTH antagonist comprises a contiguous
5 portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics:

- a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- 10 b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- c) said PTH antagonist has a minimal length of three amino acid residues.

19. A method for preventing, treating or delaying a disease or disorder
15 associated with excessive bone mineral loss in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion
20 of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics:

- a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- 25 b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- c) said PTH antagonist has a minimal length of three amino acid residues, whereby said disease or disorder associated with excessive bone mineral loss is prevented, treated or delayed.

30

20. The method of claim 19, wherein the mammal is a human.

21. The method of claim 20, wherein the human:
- a) is in need of increased bone density or bone healing;
 - b) has undergone or is presently undergoing corticosteroid therapy, chemotherapy for post menopausal bone loss, radiation therapy for cancer or hormone replacement therapy;
 - c) is immobilized or subjected to extended bed rest due to bone injury;
 - d) suffers from alcoholism, diabetes, hyperprolactinemia, anorexia nervosa, primary and secondary amenorrhea, or oophorectomy;
 - e) is 50 years or older;
 - f) is a female; or
 - g) is a male.
22. The method of claim 21, wherein the female is 40 years or older or is in the post-menopausal stage.
23. The method of claim 20, wherein the human has an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist.
24. The method of claim 20, wherein the human has an abnormal PTH agonist level.
25. The method of claim 23, wherein the abnormal comparative value between PTH agonist and PTH antagonist is determined by determining and comparing at least two of the parameters selected from the group consisting of the level of the PTH agonist, the PTH antagonist and the sum of the PTH agonist and the PTH antagonist level.
26. The method of claim 25, wherein the comparison is in the form of a ratio, a proportion or subtraction difference.

27. The method of claim 20, wherein the human has a PTH agonist/PTH antagonist ratio more than 2 and the method is used to bring the PTH agonist/PTH antagonist ratio within a range from about 1 to about 2.
- 5 28. The method of claim 19, wherein the disease or disorder associated with excessive bone mineral loss is selected from the group consisting of hyperparathyroidism, renal osteodystrophy, osteoporosis and parathyroid cancer.
- 10 29. The method of claim 19, wherein the N-terminal amino acid residue of the PTH antagonist starts at position 2 of the PTH₁₋₈₄.
30. The method of claim 19, wherein the C-terminal amino acid residue of the PTH antagonist ends at position 84 of the PTH₁₋₈₄.
- 15 31. The method of claim 19, wherein the PTH antagonist is a protein or a peptide, or a nucleic acid encoding said protein or peptide, selected from the group consisting of PTH₂₋₈₄, PTH₃₋₈₄, PTH₄₋₈₄, PTH₅₋₈₄, PTH₆₋₈₄, PTH₇₋₈₄, PTH₈₋₈₄, PTH₉₋₈₄, PTH₁₀₋₈₄, PTH₁₁₋₈₄, PTH₁₂₋₈₄, PTH₁₃₋₈₄, PTH₁₄₋₈₄, PTH₁₅₋₈₄, PTH₁₆₋₈₄, PTH₁₇₋₈₄, PTH₁₈₋₈₄, PTH₁₉₋₈₄, PTH₂₀₋₈₄, PTH₂₁₋₈₄, PTH₂₂₋₈₄, PTH₂₃₋₈₄, PTH₂₄₋₈₄, PTH₂₅₋₈₄, PTH₂₆₋₈₄, PTH₂₇₋₈₄, PTH₂₈₋₈₄, PTH₂₉₋₈₄, PTH₃₀₋₈₄, PTH₃₁₋₈₄, PTH₃₂₋₈₄, and PTH₃₃₋₈₄.
- 20 32. The method of claim 19, wherein the PTH antagonist is a protein or a peptide, or a nucleic acid encoding said protein or peptide, selected from the group consisting of PTH₇₋₆₉, PTH₇₋₇₀, PTH₇₋₇₁, PTH₇₋₇₂, PTH₇₋₇₃, PTH₇₋₇₄, PTH₇₋₇₅, PTH₇₋₇₆, PTH₇₋₇₇, PTH₇₋₇₈, PTH₇₋₇₉, PTH₇₋₈₀, PTH₇₋₈₁, PTH₇₋₈₂, PTH₇₋₈₃ and PTH₇₋₈₄.
- 25 33. The method of claim 19, wherein the PTH antagonist has a length of 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49,
- 30

50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82 or 83 amino acid residues.

34. The method of claim 19, wherein the PTH antagonist further
5 comprises an amino acid residue substitution or modification that enhances or does not decrease its antagonist activity, or an amino acid residue substitution or modification that stabilizes the PTH antagonist.

35. The method of claim 34, wherein the an amino acid residue
10 substitution or modification is selected from the group consisting of His₂₅, His₂₆, Leu₂₇, Tyr₃₄, D-Trp₁₂, Nle_{8,18}, desamino(Nle_{8,18}), Lys₁₃ modified in the epsilon-amino acid group by N,N-diisobutyl or 3-phenylpropanoyl, Gly₁₂ substituted by D-Trp, L-Trp, L- or D- α - or β -naphthylalanine, or D- or L- α -MeTrp, the amino acid residue at positions 7, 11, 23, 24, 27, 28, or 31 being cyclohexylalanine, the
15 amino acid residue at position 3, 16, 17, 18, 19, or 34 being α -aminoisobutyric acid, the amino acid residue at position 1 being α , β -diaminopropionic acid, the amino acid residue at position 27 being homoarginine, the amino acid residue at position 31 being norleucine, each of Arg₂₅, Lys₂₆, Lys₂₇ being substituted with Ala, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Met, Phe, Pro, Ser, Thr, Trp, Tyr
20 or Val, and a combination thereof.

36. The method of claim 19, wherein the PTH antagonist is administered
in a solid or a liquid dosage form.

37. The method of claim 19, wherein the PTH antagonist is administered
25 orally, parenterally, intranasally, topically, injectably or via a liposome.

38. The method of claim 19, wherein the nucleic acid encoding the PTH
antagonist is administered via a gene therapy vector.
30

39. The method of claim 19, wherein the PTH antagonist is administered as a bolus.

40. The method of claim 19, wherein the PTH antagonist is administered
5 continuously.

41. The method of claim 19, wherein the PTH antagonist is administered intermittently or is multiply administered.

42. The method of claim 41, wherein the PTH antagonist is administered
10 over a course of about 1, 2, 2-6, 6-12, or 12-24 hours.

43. The method of claim 41, wherein the PTH antagonist is administered
15 over a course of about 1, 2, 2-5, 5-14, or 14-60 days.

44. The method of claim 41, wherein the PTH antagonist is administered
over a course of about 1, 2, 2-6, 6-12, 12-24, 24-48, or more months.

45. The method of claim 41, wherein the PTH antagonist is administered
20 intraperitoneally daily for about 2 months and then the administration is stopped for about 1 month and then resumed intraperitoneally daily for about 2 months.

46. The method of claim 20, wherein the human has undergone or is
25 presently undergoing PTH agonist therapy, and the method is used to prevent, treat or delay excessive bone mineral loss caused by or associated with the PTH agonist therapy.

47. The method of claim 46, which is used to prevent, treat or delay bone
30 density decrease caused by or associated with said PTH agonist therapy.

48. The method of claim 46, wherein the PTH agonist has the PTH adenylate cyclase activating activity.

5 49. The method of claim 46, wherein the human has an abnormal PTH agonist and/or PTH antagonist level.

10 50. A method for preventing, treating or delaying the effect of a PTH agonist in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics:

- 15 a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- 20 c) said PTH antagonist has a minimal length of three amino acid residues, whereby the effect of said PTH agonist is prevented, treated or delayed.

51. The method of claim 50, wherein the PTH agonist has the PTH adenylate cyclase activating activity.

25 52. The method of claim 50, wherein the human has an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist.

30 53. The method of claim 50, wherein the human has a PTH agonist/PTH antagonist ratio more than 2 and the method is used to bring the PTH agonist/PTH antagonist ratio within a range from about 1 to about 2.

54. The method of claim 50, wherein the PTH agonist is a human PTH agonist.

55. The method of claim 54, wherein the PTH agonist comprises a
5 contiguous portion of human PTH having an amino acid sequence set forth in
SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH,
and said PTH agonist has the following characteristics:

- a) the N-terminal amino acid residue of said PTH agonist starts at
position 1 of said PTH₁₋₈₄; and
- 10 b) the C-terminal amino acid residue of said PTH agonist ends at any
position spanning position 34 through position 84 of said PTH₁₋₈₄.

56. The method of claim 50, which is used for treating hypercalcemia.

15 57. The method of claim 50, which is used for diagnosing or treating
hyperparathyroidism or pseudohypoparathyroidism.

58. The method of claim 57, wherein the hyperparathyroidism results in a
hypercalcemic crisis.

20 59. The method of claim 57, wherein the hyperparathyroidism is caused by
renal failure.

60. The method of claim 50, wherein the mammal has a tumor producing a
25 parathyroid hormone-like substance.

61. The method of claim 50, which is used for treating an immune disease.

62. The method of claim 61, wherein the immune disease comprises
30 inflammation, an allergic response or hyperactive lymphocytes.

63. The method of claim 50, which is used for treating hypertension.

64. The method of claim 50, wherein the mammal is a human patient undergoing a calcium or vitamin D treatment and the method further comprises a
5 step of monitoring the PTH antagonist level or a comparative value between PTH agonist and PTH antagonist to guide the administration of calcium or vitamin D in the patient.

65. A method for identifying a subject having or at risk of having
10 osteoporosis or decreased bone density, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist as having or at risk of having osteoporosis or decreased bone density.

15

66. The method of claim 65, wherein the abnormal PTH agonist/PTH antagonist ratio is more than 2 or less than 1.

67. A method for identifying a subject in need of parathyroid hormone
20 (PTH) antagonist treatment, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist as in need of parathyroid hormone (PTH) antagonist treatment.

25

68. The method of claim 67, wherein the abnormal PTH agonist and/or PTH antagonist level results in a PTH agonist/PTH antagonist ratio more than 2 or less than 1.

30

69. The method of claim 67, further comprising administering an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances

production and/or antagonizing function of said PTH antagonist to the identified subject.

5 70. The method of claim 69, wherein the PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics:

- a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- 10 b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- c) said PTH antagonist has a minimal length of three amino acid residues.

15 71. A method for monitoring a subject undergoing treatment for osteoporosis or decreased bone density, which method comprises determining PTH antagonist level or a comparative value between PTH agonist and PTH antagonist and identifying a subject having an abnormal PTH antagonist level or an abnormal comparative value between PTH agonist and PTH antagonist.

20 72. The method of claim 71, which identifies a subject having PTH agonist/PTH antagonist ratio more than 2 or less than 1 as having an abnormal comparative value between PTH agonist and PTH antagonist.

25 73. The method of claim 72, further comprising bringing the PTH agonist/PTH antagonist ratio within a range from about 1 to about 2.

30 74. The method of claim 71, which identifies a subject having PTH agonist-PTH antagonist equals or is more than 50 pg/ml or a subject having PTH agonist level is less than PTH antagonist level as having an abnormal comparative value between PTH agonist and PTH antagonist.

75. The method of claim 71, wherein the subject is undergoing a calcium, bisphosphanate or vitamin D treatment.

76. A method for identifying an agent suitable for preventing, treating or
5 delaying osteoporosis, which method comprises:

- a) measuring PTH antagonist activity in the presence and absence of a candidate compound; and
- b) identifying a compound that enhances said PTH antagonist activity as an agent suitable for preventing, treating or delaying osteoporosis.

10 77. The method of claim 76, wherein the PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics:

- 15 a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- c) said PTH antagonist has a minimal length of three amino acid residues.

20 78. A pharmaceutical composition, which pharmaceutical composition comprises an effective amount of a parathyroid hormone (PTH) antagonist and a pharmaceutically acceptable carrier or excipient, wherein said PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set
25 forth in SEQ ID NO:2, dog PTH having an amino acid sequence set forth in SEQ ID NO:3, bovine PTH having an amino acid sequence set forth in SEQ ID NO:4, rat PTH having an amino acid sequence set forth in SEQ ID NO:5, or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6, or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH
30 antagonist has the following characteristics:

- a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- 5 c) said PTH antagonist has a minimal length of three amino acid residues.

79. A method for preventing, treating or delaying a disease or disorder associated with excessive bone mineral loss in a mammal, which method comprises administering to a mammal, to which such prevention, treatment or
10 delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set forth in SEQ ID NO:2, dog PTH having an amino acid sequence set forth in SEQ ID NO:3, bovine PTH having an
15 amino acid sequence set forth in SEQ ID NO:4, rat PTH having an amino acid sequence set forth in SEQ ID NO:5 or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6, or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH antagonist has the following characteristics:

- 20 a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- c) said PTH antagonist has a minimal length of three amino acid residues,
25 whereby said disease or disorder associated with excessive bone mineral loss is prevented, treated or delayed.

80. A method for preventing, treating or delaying the effect of a PTH agonist in a mammal, which method comprises administering to a mammal, to
30 which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances

production and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of pig PTH having an amino acid sequence set forth in SEQ ID NO:2, dog PTH having an amino acid sequence set forth in SEQ ID NO:3, bovine PTH having an amino acid sequence set forth in
5 SEQ ID NO:4, rat PTH having an amino acid sequence set forth in SEQ ID NO:5 (PTH₁₋₈₄), or chicken PTH having an amino acid sequence set forth in SEQ ID NO:6, or a nucleic acid encoding said portion of pig, dog, bovine, rat or chicken PTH, and said PTH antagonist has the following characteristics:

- 10 a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- c) said PTH antagonist has a minimal length of three amino acid residues, whereby the effect of said PTH agonist is prevented, treated or delayed.

15

81. A method for preventing, treating or delaying bone metastasis in a human, which method comprises administering to a human, to which such prevention, treatment or delay is needed or desirable, an effective amount of a parathyroid hormone (PTH) antagonist or an agent that enhances production
20 and/or antagonizing function of said PTH antagonist, wherein said PTH antagonist comprises a contiguous portion of human PTH having an amino acid sequence set forth in SEQ ID NO:1 (PTH₁₋₈₄), or a nucleic acid encoding said portion of human PTH, and said PTH antagonist has the following characteristics:

- 25 a) the N-terminal amino acid residue of said PTH antagonist starts at any position spanning position 2 through position 33 of said PTH₁₋₈₄;
- b) the C-terminal amino acid residue of said PTH antagonist ends at any position spanning position 35 through position 84 of said PTH₁₋₈₄; and
- c) said PTH antagonist has a minimal length of three amino acid residues, whereby said bone metastasis is prevented, treated or delayed.

30

82. The method of claim 81, wherein the bone metastasis is associated with or caused by breast cancer, prostate cancer or multiple myeloma.

5 83. The method of claim 81, which is used to prevent, treat or delay morbidity or mortality associated with the bone metastasis.

84. The method of claim 81, which is used to prevent, treat or delay pain, pathological fractures, hypercalcemia or spinal cord compression associated with the bone metastasis.

10

85. The method of claim 81, wherein the PTH antagonist or an agent that enhances production and/or antagonizing function of the PTH antagonist is administered orally or intravenously.

15


86. The method of claim 81, which is used in combination with a hormone therapy or a chemotherapy.

87. A kit, which kit comprises a pharmaceutical composition of claim 1 in a container and an instruction for using the pharmaceutical composition in preventing, treating or delaying bone metastasis.

20

FIGURE 1

Whole Human PTH (1-84)



2/7

FIGURE 2





FIGURE 3



3/7





FIG. 4

4/7

FIGURE 5



5/7




FIG.6A




FIG.6B




FIG.7

6/7

FIGURE 8





FIGURE 9



7/7




FIG.10




FIG.11

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.